



Tansley review

Phosphorus acquisition and use: critical adaptations by plants for securing a nonrenewable resource

Author for correspondence:
Carroll P. Vance
Tel: +1 612 6255715
Fax: +1 651 6495058
Email: vance004@tc.umn.edu

Received: 14 May 2002
Accepted: 10 October 2002

Carroll P. Vance,^{1,2} Claudia Uhde-Stone^{2,3} and Deborah L. Allan³

¹USDA, Agricultural Research Service, Plant Science Research Unit, University of Minnesota 1991 Upper Buford Circle, St Paul, MN 55108, USA; ²Departments of Agronomy and Plant Genetics, University of Minnesota 1991 Upper Buford Circle, St Paul, MN 55108, USA, ³Soil, Water and Climate, University of Minnesota 1991 Upper Buford Circle, St Paul, MN 55108, USA

Contents

I. Introduction	424	VI. P deficiency modifies carbon metabolism	431
II. The phosphorus conundrum	424	VII. Acid phosphatase	436
III. Adaptations to low P	424	VIII. Genetic regulation of P responsive genes	437
IV. Uptake of P	424	IX. Improving P acquisition	439
V. P deficiency alters root development and function	426	X. Synopsis	440

Summary

Key words: roots, expressed sequence tags (ESTs), nutrient acquisition, white lupin (*Lupinus albus*), genetics, plant stress.

Phosphorus (P) is limiting for crop yield on > 30% of the world's arable land and, by some estimates, world resources of inexpensive P may be depleted by 2050. Improvement of P acquisition and use by plants is critical for economic, humanitarian and environmental reasons. Plants have evolved a diverse array of strategies to obtain adequate P under limiting conditions, including modifications to root architecture, carbon metabolism and membrane structure, exudation of low molecular weight organic acids, protons and enzymes, and enhanced expression of the numerous genes involved in low-P adaptation. These adaptations may be less pronounced in mycorrhizal-associated plants. The formation of cluster roots under P-stress by the nonmycorrhizal species white lupin (*Lupinus albus*), and the accompanying biochemical changes exemplify many of the plant adaptations that enhance P acquisition and use. Physiological, biochemical, and molecular studies of white lupin and other species response to P-deficiency have identified targets that may be useful for plant improvement. Genomic approaches involving identification of expressed sequence tags (ESTs) found under low-P stress may also yield target sites for plant improvement. Interdisciplinary studies uniting plant breeding, biochemistry, soil science, and genetics under the large umbrella of genomics are prerequisite for rapid progress in improving nutrient acquisition and use in plants.

© *New Phytologist* (2003) **157**: 423–447

I. Introduction

Phosphorus (P) is one of 17 essential elements required for plant growth (Bieleski, 1973; Raghothama, 1999). The P concentration in plants ranges from 0.05 to 0.50% dry weight. This element plays a role in an array of processes, including energy generation, nucleic acid synthesis, photosynthesis, glycolysis, respiration, membrane synthesis and stability, enzyme activation/inactivation, redox reactions, signaling, carbohydrate metabolism, and nitrogen (N) fixation. The concentration gradient from the soil solution P to plant cells exceeds 2000-fold, with an average inorganic phosphate (P_i) concentration of $1 \mu\text{M}$ in the soil solution (Bieleski, 1973; Schachtman *et al.*, 1998). This concentration is well below the K_m for plant uptake. Thus, although bound P is quite abundant in many soils, it is largely unavailable for uptake. As such, P is frequently the most limiting element for plant growth and development. Crop yield on 30–40% of the world's arable land is limited by P availability (Runge-Metzger, 1995; von Uexküll & Mutert, 1995). Phosphorus is unavailable because it rapidly forms insoluble complexes with cations, particularly aluminum and iron under acid conditions. The acid-weathered soils of the tropics and subtropics are particularly prone to P deficiency (Sanchez & Uehara, 1980). Application of P-containing fertilizers is usually the recommended treatment for enhancing soil P availability and stimulating crop yields.

II. The phosphorus conundrum

Application of P fertilizer, however, is problematic for both the intensive and extensive agriculture of the developed and developing worlds, respectively. In intensive agriculture, a maize crop yield of $6\text{--}9 \text{ t ha}^{-1}$ requires crop uptake of $30\text{--}50 \text{ kg P ha}^{-1}$ (Ellington, 1999; Vance, 2001; Johnston, 2002) with about two-thirds of that removed in the harvested portion of the crop. Small grains yielding 3 t ha^{-1} take up $15\text{--}22 \text{ kg P ha}^{-1}$, again with a 70% removal rate. Soybean takes up $20\text{--}25 \text{ kg P ha}^{-1}$, with 80–100% removed in the harvested portion (Johnston, 2002). However, even under adequate P fertilization, only 20% or less of that applied is removed in the first year's growth because of retention by the soil (Russell, 1973). This results in P loading of prime agricultural land. Run-off from P-loaded soils is a primary factor in eutrophication and hypoxia of lakes and marine estuaries of the developed world (Runge-Metzger, 1995; Bumb & Baanante, 1996). Another reason for alarm is that by some estimates inexpensive rock phosphate reserves could be depleted in as little as 60–80 yr (CAST, 1988; Vance, 2001). Phosphorus fertilizer use increased four- to five-fold between 1960 and 2000 and is projected to increase further by 20 Tg yr^{-1} by 2030 (Table 1). As noted by Abelson (1999), a potential phosphate crisis looms for agriculture in the twenty-first century. An even greater concern than the excessive use of

Table 1 Agriculture production and resource use in the recent past to the near future

Item	1960	2000	2030–2040
Food production (Mt)	1.8×10^9	3.5×10^9	5.5×10^9
Population (billions)	3	6	8 (perhaps 10)
Irrigated land (% of arable)	10	18	20
Cultivated land (ha)	1.3×10^9	1.5×10^9	1.8×10^9
Water-stressed countries	20	28	52
N fertilizer use (Tg)	10	88	120
P fertilizer use (Tg)	9	40	55–60

Data adapted from Vance (2001). Mt, metric tons; Tg, 10^{12} g or million metric tons.

P fertilizers by intensive agriculture, is the lack of available P fertilizers for extensive agriculture in the tropics and subtropics where the majority of Earth's people live. Lack of fertilizer infrastructure, money for purchase, and transportation make P fertilization unattainable for these areas. Sustainable management of P in agriculture requires that plant biologists discover mechanisms that enhance P acquisition and exploit these adaptations to make plants more efficient at acquiring P, develop P-efficient germplasm, and advance crop management schemes that increase soil P availability.

III. Adaptations to low P

Plants have evolved two broad strategies for P acquisition and use in nutrient-limiting environments: (1) those aimed at conservation of use; and (2) those directed toward enhanced acquisition or uptake (Lajtha & Harrison, 1995; Horst *et al.*, 2001; Vance, 2001). Processes that conserve the use of P involve decreased growth rate, increased growth per unit of P uptake, remobilization of internal P_i , modifications in carbon metabolism that bypass P-requiring steps, and alternative respiratory pathways (Schachtman *et al.*, 1998; Plaxton & Carswell, 1999; Raghothama, 1999; Uhde-Stone *et al.*, 2002a,b). By comparison, processes that lead to enhanced uptake include increased production and secretion of phosphatases, exudation of organic acids, greater root growth along with modified root architecture, expansion of root surface area by prolific development of root hairs, and enhanced expression of P_i transporters (Marschner *et al.*, 1986; Duff *et al.*, 1994; Schachtman *et al.*, 1998; Gilroy & Jones, 2000; Lynch & Brown, 2001). Although beyond the scope of this review, the most prevalent evolutionary adaptation by land plants (80% of all species) for acquiring P is through mycorrhizal symbioses (Koide & Kabir, 2000; Smith *et al.*, 2000; Burleigh *et al.*, 2002; Tibbett & Sanders, 2002).

IV. Uptake of P

Phosphorus is taken up by plants in the orthophosphate (P_i) forms H_2PO_4^- and HPO_4^{2-} , which occur in soil solutions at

very low concentrations (0.1–10 μM ; Hinsinger, 2001). A pH optimum for P_i uptake of 4.5–5.0 indicates preferential plant uptake of H_2PO_4^- over HPO_4^{2-} (Raghothama, 1999). Although total soil P content typically varies from 500 to 2000 p.p.m., total bioavailable P, as measured by soil extractants may be only a few p.p.m. (Sanyal & DeDatta, 1991). Up to half the soil P can be organic, derived from plant residues and soil organisms. Organic P must be mineralized before it can be taken up by plants (Horst *et al.*, 2001).

The reactions controlling the amounts of P_i in solution include dissolution–precipitation of P-bearing minerals, adsorption–desorption of phosphate on soil surfaces, and the hydrolysis of organic matter (Matar *et al.*, 1992; Comerford, 1998; Hinsinger, 2001). It can be difficult to distinguish adsorbed and precipitated forms of P, but retention or ‘fixation’ of P by soil components is greatest in the presence of Fe- and Al-hydroxylated surfaces (from Fe and Al oxides and clay minerals) and, at higher pH, calcium carbonate (Matar *et al.*, 1992; Comerford, 1998). Along with the types and amounts of clay and metal oxides, P availability is also controlled by soil solution pH, ionic strength, concentrations of P and metals (Fe, Al and Ca) and the presence of competing anions, including organic acids (Sanyal & DeDatta, 1991; Hinsinger, 2001). Thus, plant roots (and microbes) can alter solution P_i availability by acidification of the rhizosphere, exudation of organic acids, and secretion of extracellular phosphatases (Comerford, 1998; Hinsinger, 2001).

Because of its strong reactions with soil components, P_i is principally supplied to plant roots by diffusion rather than mass flow (Comerford, 1998; Hinsinger, 2001). At the root surface, P_i is rapidly taken up, resulting in a P_i depletion shell of 0.2–1.0 mm around the root (Barber *et al.*, 1963; Holford, 1997). Although the soil solution P_i concentration rarely exceeds 2 μM , that in plant cells is much higher, 2–20 mM (Bialeski, 1973; Schachtman *et al.*, 1998). For the plant to surmount this concentration difference as well as the negative membrane potential, active transport across the plasmalemma is required. Physiological research over the last few decades, including both inhibitor and kinetic studies, laid the foundation for the molecular investigation of transport processes (Epstein, 1953; Smith, 2001). The striking reduction in P_i accumulation of tissues treated with inhibitors reflects the energy requirement for uptake (Bialeski, 1973; Raghothama, 1999). Moreover, kinetic analysis of P_i uptake shows that plants have both a low- and high-affinity uptake system (Bialeski, 1973; Smith *et al.*, 2000). The high-affinity system operating at low P_i concentrations has an apparent K_m ranging from 3 to 10 μM ; the low-affinity system operating at high P_i concentrations has a K_m ranging from 50 to 300 μM (Nandi *et al.*, 1987; Furihata *et al.*, 1992). The high-affinity uptake process is induced when P_i is deficient whereas the low-affinity system appears to be constitutive in plants (Raghothama, 1999). Phosphorus moves symplastically from the root surface to the xylem at a rate of about 2 mm h^{-1} (Bialeski, 1973).

Xylem flow then transports P_i to the above-ground organs, where symplastic transport carries it to cells within individual tissues. The movement of P_i from the xylem to the cell cytoplasm and from cytoplasm to vacuole is also against a steep electrochemical potential gradient which requires energized transport (Ullrich & Novacky, 1990). Because most P_i is transported as H_2PO_4^- , cotransport involves a cation. Acidification of the cytoplasm which occurs upon P_i addition to P-deficient cells suggests H^+ is the cotransport product for the vast majority of plants (Schachtman *et al.*, 1998). However, authors of a recent study on internodal cells of the giant alga *Chara* have reported a Na^+ -coupled P_i uptake system that was induced by P_i deficiency (Reid *et al.*, 2000).

Within recent years, functional complementation of yeast mutants defective in P_i transport has been used to isolate and characterize P_i transporters from a diverse array of plants (Muchhal *et al.*, 1996; Leggewie *et al.*, 1997; Daram *et al.*, 1998; Liu *et al.*, 1998; Liu *et al.*, 1998). Molecular characterization of P_i transporters coupled to the discovery of as many as 16 P_i transport genes within the genome of *Arabidopsis* confirms that plants have a multiplicity of P_i transporters functional in specific organs and tissues (Raghothama, 1999; Gilroy & Jones, 2000). The deduced amino acid sequence of plant P_i transporters indicate they have an M_r of ≈ 58 kDa and are comprised of 525–550 amino acid residues. All characterized P_i transporters have 12 transmembrane domains that occur as two groups (6 + 6) connected by a hydrophobic domain of 60 amino acids (Schachtman *et al.*, 1998; Smith *et al.*, 2000). Computer modeling predicts that both *N*- and *C*-terminal domains of high-affinity P_i transporters are found on the inner membrane surface (Fig. 1), while in the low-affinity transporters these domains are on the exterior surface of the membrane. Initial P_i uptake kinetic studies with a yeast mutant complemented with a P_i transporter disappointingly showed an apparent uptake K_m to be greater than 100 μM , suggesting that it was not a high-affinity transporter (Leggewie *et al.*, 1997). This reduced affinity may also reflect that expression was in a heterologous system. However, more recent studies have yielded a K_m of 31 μM for the tomato *LePT1* P_i transporter (Daram *et al.*, 1998). An even higher affinity, apparent K_m of 3.1 μM , was shown when the *Arabidopsis* PHT1 P_i transporter gene was expressed in cultured tobacco cells (Mitsukawa *et al.*, 1997). *In situ* hybridization and immunolocalization of high-affinity P_i transporters transcripts and protein in root epidermal and root hair cells (Daram *et al.*, 1998; Liu *et al.*, 1998; Chiou *et al.*, 2001) suggests their importance for P_i uptake from soil solution. Localization of plant uptake systems in the youngest regions of roots and in the root hairs is most effective for recovery of nutrients from undepleted soil (Smith *et al.*, 2001). As a depletion zone develops, effective uptake of P_i behind the root hair zone requires either mycorrhizal uptake (through hyphae that bridge the depletion zone) or plant adaptations, such as cluster roots, that solubilize and extract additional P_i from localized soil patches.




Fig. 1 Predicted topology of the high-affinity inorganic phosphate (P_i) transporter LaPT1 from white lupin based upon homology to other high-affinity P_i transporters.

Fig. 2 Transcript abundance of phosphate (P) transporter (LaPT1) and secreted acid phosphatase (sAP1) in phosphate-sufficient and P-deficient white lupin at 14 d after emergence (DAE). Plants grown in presence of sufficient P are designated by (+) and those grown under P stress are designated by (-). Letters above each lane represent: N, normal roots; Pr, proteoid (cluster) roots; S, stems; L, leaves; F, flowers; Pd, pods. Flowers and pods did not form on -P plants. Each lane contains 20 μ g of total RNA. (Derived from Liu *et al.*, 2001 and Miller *et al.*, 2001.)

When placed in +P nutrient solution, uptake of P_i on a gram fresh weight basis by P-deficient cluster roots of white lupin is much greater than that of P-sufficient plants (Keerthisinghe *et al.*, 1998; Neumann *et al.*, 1999). Moreover, the apparent K_m for P_i uptake of cluster roots is 8.6 μ M compared with a K_m of 30.7 μ M for P-sufficient controls (Neumann *et al.*, 1999). These results suggest that a high-affinity P_i uptake system is induced in cluster roots of P-deficient plants. Liu *et al.* (2001) have recently characterized a high-affinity type P_i uptake gene (*LaPT1*) from cluster roots of white lupin (*Lupinus albus*) that shows highly intensified expression in P-deficient plants. Similar to other plant P_i transporters, the white lupin cluster root P_i transporter has a 1620 bp open reading frame that encodes a protein of 540 amino acids with a M_r of 59 kDa organized into 12 transmembrane domains (Fig. 1). The deduced amino acid sequence of *LaPT1* is 85% similar to previously reported high-affinity type P_i transporters but only 75% similar to low affinity type P_i transporters. Transcripts of *LaPT1* are highly expressed in P_i -deficient cluster roots, normal roots, and stems with little to no expression in P-sufficient plants (Fig. 2). Thus, the enhanced uptake of P_i displayed by P-deficient cluster roots may be directly related to increased expression of a high-affinity type P_i transporter.

V. P deficiency alters root development and function

Shoot and fruit development, because of their agricultural and horticultural importance, have historically received much more attention than that of roots. In recent years, with a flush of interest in nutrient acquisition, studies of root biology have become more attractive. Roots are the primary source of all the mineral elements required for plant growth. As such, root growth and development are highly plastic (Klepper, 1992; McCulley, 1995; Neumann & Martinoia, 2002) and vary greatly depending upon numerous soil factors including: nutrient availability, water content, oxygen concentration, soil density, pH and gravity (Russell, 1973; Lynch, 1995; McCulley, 1995; Charlton, 1996; Raven & Edwards, 2001). Plant roots typically respond to P deficiency through allocation of more carbon to roots, resulting in increased root growth, enhanced lateral root formation, greater exploration of the surface soil, increased length and number of root hairs (Lynch, 1995; Gilroy & Jones, 2000; Liao *et al.*, 2001; Lynch & Brown, 2001; Williamson *et al.*, 2001), enhanced expression of P transporters (Muchhal *et al.*, 1996; Leggewie *et al.*, 1997; Smith *et al.*, 1997; Liu *et al.*, 1998; Liu *et al.*,

2001), and exudation of constituents (organic acids, acid phosphatases) that increase P availability (Dinkelaker *et al.*, 1989; Duff *et al.*, 1991; Tadano & Sakai, 1991; Johnson *et al.*, 1994, 1996a; Gilbert *et al.*, 1998; Gilbert *et al.*, 1999; Neumann & Römheld, 1999).

Coordinated expression of plant genes within a given environment trigger biochemical events that give rise to plant organs and tissues (Torrey, 1986; Boerjan *et al.*, 1992; Celenza *et al.*, 1995; Lynch, 1995; Scheres & Wolkenfelt, 1998; Gilroy & Jones, 2000). Changes in either gene expression or the environment can alter the developmental fate of any given organ. Understanding the genetic and biochemical mechanisms that regulate the development of plant organs are cardinal questions in plant biology. Studies of *Arabidopsis*, common bean (*Phaseolus vulgaris*) and white lupin have proven particularly valuable for revealing genetic and biochemical factors that mediate plant adaptations to P deficiency.

Root architecture

Root architecture (Lynch, 1995; Lynch & Brown, 2001) refers to the complexity of root system spatial configurations that arise in response to soil conditions. It includes root morphology, topology and distribution patterns. Soil P limitation is a primary effector of root architecture (Charlton, 1996; Johnson *et al.*, 1996b; Borch *et al.*, 1999; Williamson *et al.*, 2001) and is known to impact on all aspects of root growth and development. The classic studies of Drew (1975) and Jackson *et al.* (1990) demonstrated the effect of localized supply of soil P on root proliferation in grass species. Elegant experiments with common bean coupled to simulation modeling have shown genotypic adaptations to P deficiency involve changes in root architecture that facilitate acquisition of P from the topsoil (Ge *et al.*, 2000; Liao *et al.*, 2001; Lynch & Brown, 2001). Adaptations that enhance acquisition of P from topsoil are important because of the relative immobility of P in soil, with the highest concentrations usually found in the topsoil and little movement of P into the lower soil profiles. Lynch and Brown (2001) refer to P-deficiency induced modifications in bean root architecture as adaptations for topsoil foraging. Root characteristics associated with improved topsoil foraging for P are a more horizontal basal-root growth angle, resulting in more shallow roots, increased adventitious root formation, enhanced lateral root formation and increased root hair density and length.

Williamson *et al.* (2001) have shown that P availability exerts a marked effect on the root system architecture of *Arabidopsis*. Growth under P-deficient conditions resulted in a redistribution of root growth from the primary root to lateral roots. Reduced primary root elongation under low P conditions was accompanied by increased lateral root density and elongation. Similar to common bean, *Arabidopsis* root biomass was concentrated near the soil surface, suggesting topsoil foraging. Moreover, *Arabidopsis* accessions with enhanced P

acquisition have the root architecture modifications mentioned previously and greater root penetration capacity (Narang *et al.*, 2000).

Root hairs

The abundant development of lateral roots associated with P-deficiency induced alterations in root architecture is almost invariably accompanied by increased root hair density and length. Root hairs are tubular-shaped cells specialized for nutrient uptake (Gilroy & Jones, 2000). They arise from root epidermal cells known as trichoblasts and undergo tip growth, thereby extending the root surface area in contact with the soil matrix (Ridge, 1995; Peterson & Farquhar, 1996). Root hairs can form as much as 77% of the root surface area of field crops (Parker *et al.*, 2000). For plants lacking mycorrhizae they are the primary site of nutrient uptake (Jungk, 2001; Schmidt, 2001). Root hair formation and growth is regulated largely by the supply of mineral nutrients, particularly NO_3^- and P (Gilroy & Jones, 2000; Jungk, 2001). In rape, spinach, and tomato, root hair length and number are inversely related to the P concentration in the plant (Jungk, 2001). Similarly, in legumes, P-deficiency results in both increased root hair density and length (Reid, 1981; Jungk *et al.*, 1990). Recent results from our laboratory show that *Medicago truncatula* responds quickly to P-deficiency with increased numbers and length of root hairs (S. Miller and C. P. Vance unpubl. data). Phosphorus uptake in barley cultivars is closely related to variation in root hair abundance and length (Gahoonia & Nielsen, 1997). In *Arabidopsis*, root hair density (Ma *et al.*, 2001) and elongation (Bates & Lynch, 2000) are regulated by P availability. Root hair density was fivefold greater in low-P than in high P-media. Ma *et al.* (2001) found that low P stimulated trichoblast number and the likelihood that trichoblasts would form root hairs. The average length of root hairs on P-deficient *Arabidopsis* was threefold greater than that on P-sufficient plants. Grierson *et al.* (2001) report that at least 40 genes in *Arabidopsis* affect root hair initiation and development. Many of these may be responsive to P-deficiency.

Cluster roots (proteoid roots)

Although much can be garnered from *Arabidopsis* as a model for analysis of the fundamental underpinnings of root development, alternative models must be invoked for determining how P deficiency affects root growth of crop and native species. Native species of Western Australia and South Africa are adapted to grow on the most heavily leached soils in the world (Pate *et al.*, 2001). Species of these regions have evolved specialized structures, cluster roots, for nutrient acquisition from impoverished soils. Beside mycorrhizal associations, cluster roots are regarded as one of the major adaptations for P acquisition (Skene, 1998; Pate & Watt,

2001). Cluster roots are formed in most members of the Proteaceae and in several other plant species adapted to habitats of extremely low soil fertility, including members of the Betulaceae, Casuarinaceae, Cucurbitaceae, Cyperaceae, Eleagnaceae, Leguminosae, Moraceae, Myricaceae and Restionaceae (Louis *et al.*, 1990; Dinkelaker *et al.*, 1995; Skene, 2000; Adams & Pate, 2002). In native habitats, many plant species that form cluster roots are slow-growing, sclerophyllous shrubs and trees that grow on severely P-deficient soils, such as highly leached sands, sandstones and laterites (Dinkelaker *et al.*, 1995; Pate *et al.*, 2001; Lambers *et al.*, 2002). Plant species that form cluster roots usually do not form mycorrhizal associations (Skene, 1998). The cluster roots of Proteaceae occur in close association with decomposing litter. This is especially true for *Banksia*, a genus including trees and shrubs that are dominant in the northern sandplains of south-west Western Australia. (Pate & Watt, 2001). *Banksia* typically forms dense mats of cluster roots beneath the litter layer, exploiting the relatively P-rich A horizon, while vertically descending tap roots provide access to groundwater (Jeschke & Pate, 1995; Pate & Watt, 2001). Growth of these root clusters is seasonal, starting after the onset of winter rain for nutrient uptake from the newly acquired litter during winter and spring (Lamont, 1982; Grierson & Attiwill, 1989; Jeschke & Pate, 1995; Pate & Watt, 2001). During winter, increased concentrations of P and other nutrients were found in xylem sap from cluster roots of *Banksia prionotes*, compared with the remainder of the lateral root system (Jeschke & Pate, 1995). This period of nutrient uptake is accompanied by nutrient storage in the trunk and leaves (Jeschke & Pate, 1995). It is followed by senescence of root clusters when the soil surface dries out during summer. At this stage, minerals stored in trunks and foliage are released for renewed shoot growth during summer (Jeschke & Pate, 1995).

Adaptation to P deficiency via formation of cluster roots is the result of a highly coordinated modification of root development and biochemistry. Cluster roots can comprise single clusters of very densely packed determinate lateral rootlets formed on a parent axis, as found in such Proteaceae as *Hakea* spp., *Leucadendron laureolum*, *Grevillea robusta* as well as in the legume white lupin (Gardner *et al.*, 1981; Lamont, 1993; Dinkelaker *et al.*, 1995; Skene *et al.*, 1996) (Fig. 3). *Banksia* species, however, form more complex compound clusters (Dinkelaker *et al.*, 1995; Pate & Watt, 2001; Lambers *et al.*, 2002). Several features of cluster root development and morphology are distinguished from that of typical dicot lateral roots. First, lateral roots are randomly initiated from the pericycle of primary roots near the zone of metaxylem differentiation (Torrey, 1986; Charlton, 1996), while cluster roots are initiated in waves (Skene, 2000). Second, lateral roots are initiated singularly opposite a protoxylem point, unlike cluster roots which are in multiples opposite every protoxylem point within the wave of differentiation (Fig. 4). Third, in typical

lateral roots, root hair development is highly regulated and occurs from a discrete number of epidermal cells (Ridge, 1995; Malamy & Benfey, 1997; Dolan, 2001), while cluster roots produce a superabundance of root hairs. The accompanying increase in root hair density of clustered rootlets results in an increase of surface area of greater than 100-fold compared with normal roots. Finally, contrasting with the indeterminate growth of lateral roots, cluster root growth is determinate, ceasing shortly after emergence (Fig. 3a). This highly synchronous developmental pattern indicates that cluster root formation is a finely tuned process. Moreover, because root pericycle cells are arrested in the G2 phase of the cell cycle (Skene, 1998, 2000), cluster root initiation must involve concerted release of multiple pericycle cells from the G2 phase in a wave-like pattern along second-order lateral roots.

While mycorrhizal hyphae increase the soil volume that is exploited by roots, cluster roots explore a comparatively small soil volume. The hairy and densely packed lateral rootlets bind tightly to trapped sand and particles of organic matter (Lamont, 1973; Grierson & Attiwill, 1989; Pate & Watt, 2001). The dense aggregates of cluster roots are thought to mobilize sparingly soluble P_i more effectively by concentrating root exudates in localized patches (Gardner *et al.*, 1983; Grierson & Attiwill, 1989). Excretion of organic acids and acid phosphatase from cluster roots has been shown for species of the Proteaceae (Grierson, 1992; Dinkelaker *et al.*, 1995; Lambers *et al.*, 2002). Detailed studies of the carboxylate exudation in *B. grandis* showed exudation of significant amounts of a range of carboxylates when plants were grown on Fe-phosphate or Al-phosphate (Lambers *et al.*, 2002). Interestingly, different carboxylate patterns were recovered from the rhizospheres in Fe-phosphate and Al-phosphate treatments, indicating that the plants perceived a difference in the chemistry of the rhizosphere environment (Lambers *et al.*, 2002). Development of cluster roots was suppressed when plants were grown with higher P supply (K-phosphate) (Lambers *et al.*, 2002).

Although the development and function of cluster roots in Proteaceae have received growing attention, most of the detailed physiological and biochemical work on the functioning of root clusters has been performed with white lupin. This well-characterized legume has proven an illuminating model system for understanding plant adaptations to low P in species lacking mycorrhizal associations (Avio *et al.*, 1990; Watt & Evans, 1999; Skene, 2000; Neumann & Martinoia, 2002). Biochemical changes that can increase the acquisition of scarce P occur within the cluster root zones of P-stressed white lupin (Neumann & Martinoia, 2002) (Fig. 3b). Copious quantities of organic acids, H⁺ and acid phosphatase (APase) are exuded to solubilize bound P from inorganic and organic complexes (Marschner *et al.*, 1986; Gerke *et al.*, 1994; Johnson *et al.*, 1994; Dinkelaker *et al.*, 1995; Neumann *et al.*, 1999; Miller *et al.*, 2001). In addition, P_i uptake within cluster




Fig. 3 Cluster roots of white lupin and biochemical mechanism occurring in cluster roots to enhance availability of soil phosphate (P). (a) Cluster roots of P-deficient white lupin at different stages of development. Secondary lateral roots having proteoid segments were collected from 5 to 22 d after shoot emergence (DAE). Note the determinate nature of cluster roots. (Derived from Johnson *et al.*, 1996b.) (b) Biochemical adaptations occurring in proteoid roots involve extrusion of protons and carboxylates into the rhizosphere via a plasmalemma H⁺ ATPase and anion channels, respectively. Release of protons and carboxylates solubilize P from unavailable bound forms. Acid phosphatases and perhaps phytases exuded from root hairs solubilize P from organic esters. Carboxylates and protons extruded from root hairs may also make organic P-esters more soluble and susceptible to acid phosphatases. Phosphorus released from insoluble ligands and organic P esters is taken up through P transporters in the plasmalemma.

root zones is much greater on a gram fresh weight basis than that of normal roots (Keerthisinghe *et al.*, 1998; Neumann *et al.*, 1999). Molecular events that give rise to the metabolic changes which increase P acquisition by cluster roots are mediated in part by enhanced expression and accumulation of transcripts encoding P_i transporters (Liu *et al.*, 2001), exuded acid phosphatase (Miller *et al.*, 2001) and enzymes of carbon metabolism (Uhde-Stone *et al.*, 2003a,b). Recently, we have taken a functional genomics approach to investigating adaptation to P deficiency in white lupin by sequencing some 2000 expressed sequence tags (ESTs) from P-deficiency induced cluster roots (Uhde-Stone *et al.*, 2003a,b). This approach has provided us with access to several white lupin genes important for adaptation and acclimatization to low P conditions (noted in later sections).

Hormones and root architecture

As might be expected, the internal balance of the plant growth regulators auxin, ethylene, and cytokinin is thought to play a role in P-deficiency induced alterations in lateral root development and architecture, root hair formation and cluster root development. Substantial support for the role of auxin in lateral root development is derived from evidence that: (1) exogenous auxin applications stimulate lateral root formation in many species (Torrey, 1986; Blakely *et al.*, 1988; Muday & Haworth, 1994; Charlton, 1996); (2) auxin transport inhibitors block lateral root formation and this block can be alleviated by exogenous auxin (Torrey, 1986; Muday & Haworth, 1994; Reed *et al.*, 1998); (3) *Arabidopsis* mutants that overproduce auxin have enhanced lateral root formation




Fig. 4 Initiation and emergence of cluster roots from secondary lateral roots of P-deficient white lupin. Sections (2 cm) of secondary lateral roots representing the proteoid transition zone were collected at 5, 6, 7, and 8 d after shoot emergence (DAE). Root segments were cleared with sodium hypochlorite and stained with methylene blue. Bar, 1 mm. (Derived from Johnson *et al.*, 1996b.)

(Boerjan *et al.*, 1995; Celenza *et al.*, 1995; King *et al.*, 1995; Scheres & Wolkenfelt, 1998), while mutants insensitive to auxin have impaired lateral root development (Celenza *et al.*, 1995; Hobbie & Estelle, 1995; Rogg *et al.*, 2001; Fukaki *et al.*, 2002); and (4) mutants in auxin transport have altered lateral root formation (Ruegger *et al.*, 1997; Casimiro *et al.*, 2001; Marchant *et al.*, 2002). Because cluster root formation during P deficiency reflects a striking change in root development, auxins have been implicated in their formation (Lamont *et al.*, 1984; Dinkelaker *et al.*, 1995). We (Gilbert *et al.*, 2000) and others (Skene & James, 2000) have shown that exogenous addition of auxin to P-sufficient white lupin mimics cluster root formation as seen under P-deficient conditions. Moreover, Gilbert *et al.* (2000) demonstrated that auxin transport inhibitors added to P-deficient plants dramatically reduced the formation of cluster roots. The data suggest the cluster root response to P deficiency in white lupin is directly under control of auxin transport. Conversely, auxin insensitive mutants of *Arabidopsis* (*aux1*, *axr1* and *axr4*) which have reduced lateral root formation, showed typically enhanced production of laterals on low P media (Williamson *et al.*, 2001). The authors interpret the data to mean that auxin is not directly involved in the induction of lateral roots under low P. Unfortunately, auxin transport inhibitors were not added to the P-stressed mutants, neither was auxin measured; thus interpretation of this work is open to question. An auxin transport inhibitor experiment with *aux1* under P-stress would be particularly informative since *aux1* may act as an auxin transporter (Scheres & Wolkenfelt, 1998).

Although the role of ethylene in lateral root formation is less clear there is convincing evidence that ethylene plays a role in root hair formation and abundance (Kieber, 1997; Michael, 2001). Treatments that inhibit ethylene synthesis inhibit root hair formation and growth, while the stimulation of ethylene synthesis results in an increase in root hair density

and length (Tanimoto *et al.*, 1995; Masucci & Schiefelbein, 1996; Schmidt, 2001). Of the more than 40 genes affecting root hair development (Grierson *et al.*, 2001) at least eight act directly or indirectly on ethylene biosynthesis. Mutations in the ethylene signaling pathway involving the constitutive triple response *ctr1* gene cause plants to respond as though ethylene is continually present and root hair formation as well as density is much greater than wild type. Ethylene overproduction mutants *eto1*, *2*, *3* have much longer and more root hairs than normal, while the ethylene receptor mutant *etr1* has much shorter than normal root hairs. The ethylene-overproducing mutant *eto3* exhibits a root hair phenotype reminiscent of P-deficiency induced conditions (Schmidt, 2001). Data from auxin insensitive mutants *axr2*, *iaa7*, and *aux1* show root hairs that are reduced in numbers or absent, suggesting that ethylene and auxin are both involved in regulating root hair numbers.

Nutrient stress, particularly P and Fe, are known to stimulate ethylene production (Romera & Alcantara, 1994; Borch *et al.*, 1999; Waters & Blevins, 2000; Lynch & Brown, 2001). Schmidt (2001) has proposed that the mechanism of P-deficiency induced changes in root hair density and length involves ethylene through two functionally redundant pathways. Similarly, Cho & Cosgrove (2002) present evidence that root hair initiation and growth are modulated by convergent developmental and environmental pathways, with ethylene/auxin signalling a predominant feature of the environmental stress pathway.

The changes in root development and root hair formation in cluster roots of white lupin are postulated to occur through ethylene signaling (Gilbert *et al.*, 1997; Watt & Evans, 1999). Gilbert *et al.* (2000) found a significant increase in ethylene synthesis in white lupin cluster roots. Surprisingly, application of ethylene synthesis inhibitors had little to no effect on cluster roots, but root hair number and length were not

evaluated. In our EST analysis of genes expressed in P-deficient cluster roots we found several involved in ethylene biosynthesis that were highly redundant, including ACC (1-aminocyclopropane-1-carboxylate) oxidase, methionine synthase and S-adenosylmethionine synthetase (Uhd Stone *et al.*, 2003b), but whether ethylene is a primary factor in cluster root formation and function remains to be established.

The role of cytokinins in root growth and architecture, like that of ethylene, is not resolved. Traditionally, cytokinins are thought to inhibit root growth while stimulating shoot growth (Skoog & Miller, 1957). Moreover, low cytokinin/auxin ratios favor root initiation and development in plant tissue culture experiments (Christianson & Warnick, 1985). Phosphate starvation and N deficiency appear to result in decreased cytokinin content. In addition, exogenously applied cytokinin can counteract lateral root growth stimulation induced by nutrient deficiency (Salama & Wareing, 1979; Horgan & Wareing, 1980; Kuiper *et al.*, 1988) and can repress expression of several P_i-starvation responsive genes in *Arabidopsis* (Martin *et al.*, 2000). Cytokinin inhibition of root growth may also be coupled to ethylene action (Cary *et al.*, 1995). In recent years, *Arabidopsis* mutants altered in cytokinin sensitivity (Baskin *et al.*, 1995; Faure *et al.*, 1998; Cary *et al.*, 2001) have been isolated but these mutants have no clear-cut root phenotype. Neumann *et al.* (2000) have shown that exogenous application of kinetin inhibits cluster root formation in P-deficient white lupin. Also, kinetin content was increased in cluster roots compared with normal roots. While we have not measured cytokinin profiles, it is revealing that an abundant EST from cluster roots showing enhanced expression (three- to five-fold) encoded a deduced protein having high similarity to cytokinin oxidase ($E = 10^{-20}$). Cytokinin oxidase is the key enzyme implicated in cytokinin degradation (Morris *et al.*, 1999). Enhanced degradation of cytokinins in cluster root formation and/or development might be expected because low cytokinin levels favor root growth, and P deficiency reportedly results in reduced xylem sap cytokinin levels (Wagner & Beck, 1993; Binns, 1994; Martin *et al.*, 2000). Alternatively, *in planta* regulation of the potentially large quantities of cytokinins that could be released by the mass induction of cluster root meristems may require cytokinin oxidase (Morris *et al.*, 1999; Mok & Mok, 2001).

VI. P deficiency modifies carbon metabolism

Organic acid efflux

Under normal growth and development conditions, plant roots exude a wide variety of organic compounds including: simple sugars, organic acids, amino acids, phenolics, quinones (iso)-flavonoids, growth hormones, proteins, and polysaccharides (Curl & Truelove, 1986; Marschner, 1995). Exudation of organic compounds from roots can alter rhizosphere

chemistry, soil microbial populations, competition, and plant growth. Exuded compounds are functionally diverse and can be involved in a wide array of processes ranging from signaling in plant-microbe interactions, to allelopathy and nutrient acquisition (Curl & Truelove, 1986; Marschner *et al.*, 1986; Harrison, 1997).

Under the influence of environmental stress the complement of compounds found in exudates can be significantly altered in either or both quality and quantity (Marschner, 1995; Ryan *et al.*, 2001; Neumann & Martinoia, 2002). For example, when plants are attacked by soil pathogens, the amount and type of phenolics and phytoalexins are altered, depending upon the pathogen. Similarly, rhizobia bacteria and mycorrhizal fungi induce changes in amount and types of compounds released from roots. During nutritional stress (i.e. P limitation), Al toxicity, low Fe availability, and exposure to heavy metals, roots show enhanced synthesis and exudation of several organic acids (anions) (Dinkelaker *et al.*, 1989; Delhaize *et al.*, 1993; Delhaize & Ryan, 1995; Ryan *et al.*, 1995a,b, 1997; Larsen *et al.*, 1998; Neumann *et al.*, 2000). Of the changes in response to plant nutrition, those related to low P and excess Al are among those that have been most thoroughly documented. Convincing evidence now exists for exudation of malate and citrate as a principal mechanism in alleviating the edaphic stress of P-deficiency and Al-toxicity. The release of organic acids allows for the chelation of Al³⁺, Fe³⁺ and Ca²⁺, and subsequent displacement of P_i from bound or precipitated forms (Gerke, 1994; Jones, 1998; Hinsinger, 2001; Ryan *et al.*, 2001), and may also cause organic P to become more susceptible to hydrolysis by acid phosphatases (Gerke, 1994; Braum & Helmke, 1995) (Fig. 3B). In Al-toxic soils root exudation of organic acids protects plants by chelating Al³⁺ ions in the rhizosphere, thus excluding their entry into the root. Because several recent excellent reviews (Jones, 1998; Lopez-Bucio *et al.*, 2000b; Hinsinger, 2001; Ryan *et al.*, 2001) have critically evaluated organic acid synthesis and exudation from roots, we will focus on the salient features related to the role of organic acids in alleviating P-deficiency in cluster roots of white lupin, including recent evidence for a potential efflux mechanism.

When white lupin is P deficient, cluster roots synthesize and exude striking amounts of malate and citrate (Gardner *et al.*, 1982; Dinkelaker *et al.*, 1989; Johnson *et al.*, 1994; Keerthisinghe *et al.*, 1998; Neumann *et al.*, 1999). Grown under P-deficiency, cluster roots exude 20- to 40-fold more citrate and malate than P-sufficient roots (Fig. 3b). The amount of carbon exuded in these two compounds can range from 10% to greater than 25% of the total plant dry weight. Surprisingly, P-deficient plants do not appear to suffer any loss in either dry matter accumulation or N fixation until the reproductive stage of growth (Dinkelaker *et al.*, 1989; J. Schulze, G. Gilbert, C. Vance, unpubl. data). Although the internal concentrations of organic acids increase in P-deficient roots, it is not proportional to the amount released as exudates.

This suggests specific selective synthesis and exudation of malate and citrate in response to P-stress.

Measurements of CO₂ fixation and pulse-chase labeling with ¹⁴CO₂ show the carbon required for enhanced organic acid exudation from cluster roots of P-deficient white lupin is derived from both photosynthetic CO₂ fixation and dark CO₂ fixation by cluster roots (Johnson *et al.*, 1994, 1996a). Photosynthetic CO₂ fixation provides about 65% of the carbon exuded, while cluster root dark CO₂ fixation provides about 35%. Rates of dark CO₂ fixation in cluster roots are similar to those occurring in legume root nodules (Johnson *et al.*, 1996a). The increased cluster root *in vivo* CO₂ fixation is accompanied by enhanced specific activity of phosphoenolpyruvate carboxylase (PEPC), malate dehydrogenase (MDH), and citrate synthase (CS) (Johnson *et al.*, 1996a,b; Gilbert *et al.*, 1999; Uhde-Stone *et al.*, 2003a).

Similar increases in PEPC activity in response to P deficiency have been noted in chickpea (*Cicer arietinum*), oilseed rape, and *Sesbania rostrata* (Hoffland *et al.*, 1992; Moraes & Plaxton, 2000; Aono *et al.*, 2001). Steady-state amounts of PEPC, MDH and CS mRNAs are much higher in P-deficient cluster roots compared with P-sufficient roots (Johnson *et al.*, 1996a; Uhde-Stone *et al.*, 2003a) suggesting a coordinated induction of genes involved in both developmental as well as biochemical changes.

Enhanced exudation of organic acids may not only occur because of upregulation of selected reactions leading to greater synthesis but also may be due to either reduced degradation or utilization of citrate. The activity of aconitase (AC) is reduced in cluster roots (Neumann & Römheld, 1999; Neumann *et al.*, 1999). Similarly, respiration rates are also reduced in P-deficient cluster roots (Johnson *et al.*, 1994; Neumann *et al.*, 1999). Both phenomena could result in more citrate available for exudation.

Organic acid anion (OA⁻) exudation from cluster roots of white lupin coincides with a marked acidification of the rhizosphere (Marschner *et al.*, 1987; Dinkelaker *et al.*, 1989; Neumann & Martinoia, 2002). This is consistent with reports of a correlation of OA⁻ exudation and the release of protons from white lupin and chickpea roots (Neumann & Römheld, 1999; Sas *et al.*, 2001). Sas *et al.* (2001) have shown that extrusion of protons and organic acids in white lupin was highly dependent upon P supply. On an equimolar basis, H⁺ extrusion in P-deficient plants was two- to three-fold greater than organic acid exudation. Similarly, a recent report by Yan *et al.* (2002) showed substantially enhanced proton release from cluster roots of P-deficient white lupin. Proton release occurred primarily from young clusters and decreased significantly as cluster roots aged. During prolonged P deficiency in white lupin plants, OA⁻ exudation from cluster roots increased while the release of protons decreased, indicating that different mechanisms are involved in proton release and OA⁻ exudation (Sas *et al.*, 2001). A comparison with tomato (*Lycopersicon esculentum*) showed an increase of

the net release of protons from roots of P-starved tomato while OA⁻ excretion from tomato roots decreased under P deficiency (Neumann & Römheld, 1999). Wheat (*Triticum aestivum*) displayed a decrease in OA⁻ exudation under P deficiency and no proton extrusion (Neumann & Römheld, 1999).

Proton release into the rhizosphere is a common response of plant cells to counter intracellular acidity (Raven & Smith, 1976). Generally, proton release results from the activity of a plasma membrane H⁺ ATPase. This enzyme uses ATP to pump protons out of the cell, thereby creating pH and electrical potential differences across the plasmalemma. Yan *et al.* (2002) reported enhanced catalytic activity of a plasma membrane H⁺ ATPase of P-deficient cluster roots in white lupin that might be responsible for the increase in H⁺ extrusion. Characterization of the H⁺ ATPase from P-deficient cluster roots showed a threefold increase in V_{max} and a fourfold increase in H⁺ ATPase enzyme protein compared with P-sufficient roots. Only the youngest P-deficient cluster roots had significant proton extrusion and H⁺ ATPase activity. It is noteworthy that Sakano (2001) addresses the potentially problematic generation of protons during OA⁻ synthesis via glycolysis in his revised pH-stat hypothesis of the plant cell. By contrast to the 'feed-forward' fashion of the nonplant system, plant glycolysis is regulated by a feedback process. This feedback regulation only permits glycolytic activity when the cytoplasm is alkaline enough to stimulate PEPC, an enzyme with an alkaline pH optimum (Sakano, 2001). As Sakano points out, active H⁺ extrusion is a possible response of the plant cell to prevent the inhibiting effect of cytosolic acidification.

While neither the specific cell types for nor the mechanism of organic acid exudation have been characterized for cluster roots of white lupin, studies on Al tolerance in wheat (Ryan *et al.*, 1997) and maize (*Zea mays*) (Kollmeier *et al.*, 2001; Pineros & Kochian, 2001) have identified the root apex as the zone of organic acid exudation. Moreover, these patch-clamp and inhibitor studies indicate that Al may activate a malate/citrate-permeable anion channel to facilitate the exudation of large quantities of organic acids. Zhang *et al.* (2001) showed that anion channel frequency and rapidity of activation was greater in an Al³⁺-tolerant wheat line compared with a near-isogenic Al³⁺-sensitive line. Neumann *et al.* (1999) showed that exposure of P deficiency induced cluster roots to the anion channel-blockers ethacrynic- and anthracene-9-carboxylic acids inhibited citrate exudation by 40–60%. Moreover, we have recently shown through *in situ* hybridization (Uhde-Stone *et al.*, 2003a) that PEPC and MDH transcripts are localized to cluster root apices and elongation zones, indicating the enzymes necessary for malate and perhaps citrate synthesis occur in the root tips (Fig. 5). Our data coupled with that of Neumann *et al.* (1999) suggest that P deficiency induced rapid exudation of organic acids, similar to Al³⁺-tolerance, may involve selective anion channel proteins at sites of exudation. Since malate and citrate are fully dissociated in the




Fig. 5 *In situ* localization of phosphoenolpyruvate carboxylase (PEPC) in P-deficient cluster roots of white lupin. Sections A–J were hybridized with ^{35}S -labeled antisense RNA derived from LaPEPC1 cDNA, whereas section K was hybridized with the corresponding sense RNA probe. The dark-field (a, c, e, g, i and k) and bright-field (b, d, f and j) images correspond to longitudinal and transverse sections through cluster rootlets 9 d after emergence (DAE) (a, b) and 12 DAE (c–h) as well as through 12 DAE +P normal roots (i, j, k). Arrows in d indicate the approximate location of the two transverse sections shown in e–h. The areas that are white in a, c, e, g and i indicate the presence of PEPC mRNA. Sense control (K) showed no reaction. Bars are 50 μm (e, g), 100 μm (a) and 200 μm (c, i, k); pi, proteoid initials; vb, vascular bundle; c, cortex; e, epidermis, am, apical meristem, rc, rootcap. (Derived from Uhde-Stone *et al.*, 2003a.)

cytosol and because cell membranes are in essence impermeable to ions, it is not surprising that the release of organic acids from root tips in either P deficiency or Al toxicity might involve some type of channel protein.

Although a large number of cation channels and transporters belonging to a multiplicity of gene families have been identified over the past decade (Cao *et al.*, 1995; Maathuis & Sanders, 1999; Barbier-Brygoo *et al.*, 2000; Howitt & Udvardi, 2000; Theodoulou, 2000; Maser *et al.*, 2001), the molecular events involved in organic acid (anion) excretion remain elusive. While anion channels are probably ubiquitous in plants, few have been characterized (Schroeder, 1995; Barbier-Brygoo *et al.*, 2000). Mechanisms of anion transport potentially could include chloride channels (CLC), aquaporins, and multidrug extrusion proteins (MATE). To date, only putative anion channel genes related to the CLC family have been characterized (Hechenberger *et al.*, 1996; Lurin *et al.*, 1996; Geelen *et al.*, 2000). These genes encode messages of approximately 2.4 kb giving rise to proteins of approximately 785 amino acids with M_r of *c.* 85 kDa. The *Arabidopsis* *CLC-d* gene could functionally substitute for a yeast CLC protein involved in iron-limited growth, while the *CLC-a* gene appeared to be related to nitrate acquisition. Anion channel genes encoding transmembrane proteins involved in organic acid efflux have yet to be characterized.

During our annotation of ESTs from cluster roots we found that several proteins with putative transport-related functions were highly expressed in P-stressed conditions. One of these ESTs was redundant (found 10 times) and showed a ninefold induction in -P cluster roots (Uhde-Stone *et al.*, 2003b). This EST displayed high homology to a putative integral membrane protein in *Arabidopsis* belonging to the MATE protein family. MATE proteins are a large family of putative antiporters that are thought to be involved in excretion of a variety of drugs and toxins (Brown *et al.*, 1999; Debeaujon *et al.*, 2001; Diener *et al.*, 2001). Around 55 putative members of the MATE family have been predicted in *Arabidopsis*, two distantly related MATE proteins have been predicted in yeast. To date, only a few MATE proteins have been cloned (Debeaujon *et al.*, 2001; Diener *et al.*, 2001; Rogers & Guerinot, 2002). An *Arabidopsis* MATE has been shown to control response to iron deficiency. It represents either an Fe sensor or transporter of small organic molecules (Rogers & Guerinot, 2002). Analysis of the white lupin P-stress induced MATE revealed a 2.1-kb cDNA encoding a deduced amino acid sequence of 531 amino acids having 10–12 transmembrane helices, depending upon the program used for analysis. These findings make this putative MATE EST an extremely interesting candidate for a role in organic acid excretion. Functional studies in yeast and *Xenopus* oocytes will

allow us to examine the role of the MATE type protein in organic acid excretion.

Alternative glycolytic pathways

Besides increased acquisition of soil P, conservation of internal pools of acquired P_i is considered an important adaptation for growth on low P. As several enzymes of the glycolytic pathway depend on P_i or adenylates as cosubstrates, metabolic processes may be impaired by severe P deprivation. Yet the generation of energy and the production of carbon skeletons continues during conditions of P limitation, as evident in the excretion of significant amounts of organic acids. In response to limited P, many plants show remarkable flexibility in adjusting metabolic rates and utilizing alternative metabolic pathways. Alternative glycolytic reactions can bypass P_i - or ATP-requiring steps of glycolysis under environmental stress conditions such as P_i starvation or anoxia (Duff *et al.*, 1989b; Mertens, 1991; Theodorou *et al.*, 1992; Theodorou & Plaxton, 1996). While ATP and ADP-levels have been reported to decline under P deficiency (Ashihara *et al.*, 1988; Duff *et al.*, 1989b), pyrophosphate (PP_i) concentrations appear to be buffered during P_i stress (Duff *et al.*, 1989b; Dancer *et al.*, 1990). Despite an earlier assumption of cellular bioenergetics that PP_i anhydride bonds were not utilized in cellular metabolism, studies have shown that under certain environmental stresses PP_i can serve as an energy donor in the plant cytosol (Duff *et al.*, 1989b; Plaxton, 1996; Plaxton & Carswell, 1999). A well-documented alternative glycolytic pathway is catalysed by a PP_i -dependent phosphofructokinase (PFP) that, under P deficiency, can bypass the ATP-dependent phosphofructokinase (PFK), generating fructose 1,6-bisphosphate (Theodorou *et al.*, 1992; Theodorou & Plaxton, 1996; Plaxton & Carswell, 1999). Other processes that might use PP_i are the cleavage of sucrose by a PP_i -dependent sucrose synthase pathway and the active transport of protons into the vacuole by a PP_i -dependent H^+ pump in the tonoplast (Plaxton & Carswell, 1999). PP_i has been assumed to be a byproduct of secondary metabolism and anabolism, therefore use of PP_i as an energy donor helps to conserve limited ATP pools (Duff *et al.*, 1989b; Theodorou *et al.*, 1992; Plaxton, 1996; Theodorou & Plaxton, 1996; Plaxton & Carswell, 1999).

Another alternative glycolytic pathway known in plants is catalysed by the action of a nonphosphorylating NADP-dependent glyceraldehyde-3P dehydrogenase (NADP-G3PDH) that bypasses P_i -dependent NAD-G3PDH and phosphoglycerate kinase (Duff *et al.*, 1989b; Theodorou *et al.*, 1992; Plaxton & Carswell, 1999). A third bypass of the glycolytic pathway can be catalysed by the combined activities of PEPC, MDH and NAD-malic enzyme (Theodorou & Plaxton, 1993). Phosphorus stress can severely limit the activity of pyruvate kinase (PK), an enzyme requiring P_i and ADP. The PEPC, MDH and NAD-malic enzymes can bypass PK

and thus maintain the flow of carbon from glycolysis to the TCA cycle by avoiding the use of ADP but generating free P_i (Theodorou *et al.*, 1992; Plaxton & Carswell, 1999).


The described metabolic adaptations to P_i stress appear to be plant specific. P-deficiency induced PFP activity was found in plants that are adapted to infertile soils and contain very low PFP activity when grown under sufficient P, e.g. *Arabidopsis*, rape (*Brassica napus*), sugar beet (*Beta vulgaris*), and buckwheat (*Fagopyrum esculentum*) (reviewed in Plaxton & Carswell, 1999). Murley *et al.* (1998) reported that P-stress inducible PFP activity is not found in plants that form symbiotic associations with mycorrhizal fungi. In plants that typically contain abundant PFP activity when grown under sufficient P, PFP activity stayed unchanged or declined under P deficiency (in tobacco (*Nicotiana tabacum*) seedlings, bean seedlings and tomato root cultures; Plaxton & Carswell, 1999). Analysis of ESTs from P-deficient cluster roots of white lupin indicated the induction of genes encoding for PFP and the PEPC/MDH bypass of PK, but did not reveal evidence for the involvement of a nonphosphorylating NADP-dependent G3PDH (Fig. 6).

Alternative mitochondrial respiration

Phosphorus limitation impairs mitochondrial respiration in many plants. Considering the adaptive flexibility of plant responses to P deficiency that is displayed in the glycolytic pathway, adaptations which allow alternative mitochondrial electron transport under P starvation have been hypothesized (Rychter & Mikulska, 1990; Theodorou & Plaxton, 1993).

The significant reduction in cellular ADP and P_i levels in plants undergoing P limitation is thought to inhibit the cytochrome (Cyt) pathway of electron transport. Low ADP and P_i can result in a high ATP : ADP ratio which inhibits respiration, a phenomenon well known as adenylate control of respiration (Bryce *et al.*, 1990). Nonphosphorylative pathways that can bypass energy-requiring sites include rotenone-insensitive NADH dehydrogenase (Rasmusson & Moller, 1991) and the alternative oxidase (AOX) (Lambers, 1985; Lance *et al.*, 1985; Day & Wiskich, 1995). The cyanide-resistant AOX catalyses the oxidation of ubiquinol and the reduction of oxygen to water but bypasses the last two energy-conserving sites normally associated with the Cyt pathway. Alternative oxidase activity is subject to regulation by a sulfhydryl-disulfide redox system (Siedow & Umbach, 2000) and by allosteric activation via α -keto acids, especially pyruvate (Millar *et al.*, 1993; Hoefnagel *et al.*, 1995). It was proposed that induction of AOX during P deficiency might be a consequence of elevated PEPC activity.

In roots of *P. vulgaris*, P deficiency led to decreased activity of the respiratory cytochrome pathway, while alternative, cyanide-resistant respiration and the NADH : NAD ratio increased (Rychter & Mikulska, 1990; Rychter *et al.*, 1992). Indication for a possible link between AOX respiration and



P starvation has also been found in *Catharanthus roseus* (Hoefnagel *et al.*, 1993), *Chlamydomonas reinhardtii* (Weger & Dasgupta, 1993) and tobacco cell cultures (Parsons *et al.*, 1999). However, oxidative respiration rates under P deficiency can vary greatly among different plants. Respiratory O₂ consumption in the green alga *Scenedesmus obtusiusculus* was stimulated during the first 96 h of P starvation (Tillberg & Rowley, 1989), while pea (*Pisum sativum*) roots displayed no effect of P starvation on respiration (reviewed by Theodorou & Plaxton, 1996). The respiration rates of *P. vulgaris*, *Selenastrum minutum*, *Lemna gibba* and suspension-cultured *C. roseus*, however, declined under P deprivation (Theodorou & Plaxton, 1996). In P-deficient cluster roots of white lupin, up to 60% reduction in mitochondrial respiration, as measured by O₂ consumption, has been reported (Johnson *et al.*, 1994; Neumann *et al.*, 1999), accompanied by an increased NADH : NAD ratio. These results indicate that in some plants inhibition of the CN-sensitive cytochrome pathway of respiration might not be fully compensated by AOX activity.

Enhanced glycolysis in P-deficient white lupin was accompanied by significant induction of genes encoding for an alcohol dehydrogenase (Massonneau *et al.*, 2001) and a formate dehydrogenase (fdh) (Uhde-Stone *et al.*, 2003b), indicating a

possible role of fermentative processes during P deficiency. In the green alga *C. reinhardtii*, phosphate starvation resulted in the accumulation of an mRNA with homology to pyruvate formate-lyase, another enzyme of fermentative metabolism (Dumont *et al.*, 1993). Taken together, these findings indicate that the increase of glycolytic activity and the resulting organic acid synthesis under P deprivation in white lupin could be, in part, a consequence of severely impaired oxidative respiration.

Secondary metabolism

A switch from primary to secondary metabolism is a common response of plants undergoing nutrient limitation. Phosphorus deficiency typically results in the accumulation of secondary metabolites such as flavonoids and indole alkaloids (reviewed in Plaxton & Carswell, 1999). In rice culture, accumulation of the polyamine putrescine is thought to mediate growth inhibition in response to P depletion (Shih & Kao, 1996). Yamakawa *et al.* (1983) reported that P depletion induced anthocyanin accumulation in *Vitis* cell culture, while P supplementation induced primary metabolism and inhibited anthocyanin synthesis. Increased synthesis of anthocyanins is a frequent response of plants to P deficiency

and presumably functions to ameliorate photoinhibitory damage to chloroplasts (Takahashi *et al.*, 1991).

In addition to the possible involvement of phenolics in senescence, phenolic compounds can be exuded into the rhizosphere in response to P starvation and might act as chelators and/or reductants increasing the release of bound P_i . Examples of phenolics that are exuded into the rhizosphere are piscidic acid from roots of pigeon pea (*Cajanus cajan*) (Ae *et al.*, 1990), alfafuran from alfalfa (*Medicago sativa*) roots (Masaoka *et al.*, 1993), and isoflavonoids from P-deficient cluster roots of white lupin (Neumann *et al.*, 2000).

Sclerophylly, characteristic of many of the Proteaceae, is also thought to be an adaptation involving enhanced phenolic metabolism by many species that occupy low P habitats. Sclerophylls are typically considered to be hard and stiff leaves that are tough (Turner, 1994). They are characterized by a thick cuticle and outer epidermal cell wall with abundant sclerification (Loveless, 1961, 1962). Bundle sheaths and leaf margins in particular show sclerification. The secondary wall of sclerids are heavily lignified, hence the relationship to phenolic secondary metabolism. Loveless (1962) postulated that sclerophylly is an adaptation to conserve nutrients particularly P and N. Turner (1994) concluded that sclerophylly is also related to protection against herbivory.

Enzymes in the phenolic and flavonoid pathways have been reported to be upregulated in P_i -deficient plants (Plaxton & Carswell, 1999). Several white lupin ESTs with high similarity to enzymes involved in secondary metabolism were represented in our cDNA library derived from mature cluster roots of P-deficient plants, including DAHP (3-deoxy-D-arabino-heptulosonate-7-phosphate synthase), phenylalanine ammonia-lyase, chalcone synthase, chalcone isomerase, isoliquiritigenin 2'-O-methyltransferase, 4-coumarate-CoA ligase, caffeoyl-CoA O-methyltransferase, cinnamoyl-CoA reductase and cinnamyl alcohol dehydrogenase (<http://www.home.earthlink.net/~whitelupinacclimation>). Whether these enzymes are involved in either anthocyanin or lignin biosynthesis remains to be established. By contrast to primary metabolism, secondary metabolism in general does not consume as much P_i but can serve to recycle significant amounts of P_i from phosphate esters (Plaxton & Carswell, 1999; Sakano, 2001). However, secondary metabolism produces an excess of reducing equivalents. The resulting cytosolic acidification could lead to the activation of AOX and other pathways that can relieve the accumulation of reducing equivalents (Sakano, 2001).

Modified thylakoid membranes

Adaptations to P deficiency related to carbon metabolism also include modification of the lipid composition of photosynthetic membranes (Essigmann *et al.*, 1998; Yu *et al.*, 2002). The lipid composition of thylakoids is comprised primarily of: monogalactosyl diacylglycerol, digalactosyl diacylglycerol,

phosphatidyl-glycerol and sulfoquinovosyl diacylglycerol. Under P-deficiency, thylakoid membranes display decreased amounts of phospholipids accompanied by large increases in the sulfolipid, sulfoquinovosyl diacylglycerol (Benning, 1998; Yu *et al.*, 2002). *Arabidopsis* genes encoding the two critical steps in sulfolipid synthesis have recently been isolated and characterized (Essigmann *et al.*, 1998; Yu *et al.*, 2002). The first step is catalysed by SQD, a protein that converts UDP-glucose and sulfite to UDP-sulfoquinovose. The second enzyme required for sulfolipid synthesis is SQD2 which catalyses the condensation of UDP-sulfoquinovose and diacylglycerol to sulfoquinovosyl diacylglycerol (Benning, 1998). Both *SQD1* and *SQD2* have greatly enhanced expression under P-deficient conditions. Moreover, an *Arabidopsis* line with a mutation in *SQD2* is impaired in phosphate-limited growth (Yu *et al.*, 2002). Essigmann *et al.* (1998) proposed that the function of the changes in thylakoid membrane sulfolipid : phospholipid ratio would be to conserve P while maintaining membrane integrity. Anionic sulfolipid could substitute for anionic phosphatidylglycerol, thus reducing the need for membrane-bound P, yet allowing photosynthesis to continue under P-limited conditions.

VII. Acid phosphatase

A universal response by plants to P-deficiency is the synthesis of acid phosphatase enzymes (APases). These enzymes are ubiquitous in plant organs and APase activity can be detected throughout development (Duff *et al.*, 1994). APases are implicated in: providing P during seed germination from stored phytate (Biswas & Cundiff, 1991; Brinch-Pedersen *et al.*, 2002); internal remobilization of P (Duff *et al.*, 1991; del Pozo *et al.*, 1999; Baldwin *et al.*, 2001); release of P from soil organic P-esters by exudation of enzymes into the rhizosphere (Lefebvre *et al.*, 1990; Goldstein *et al.*, 1988; Miller *et al.*, 2001); and the synthesis of glycolate from P-glycolate (Christeller & Tolbert, 1978) as well as glycerate from 3-PGA during photorespiration (Randall *et al.*, 1971). Characteristically, phytases and root-secreted APase have little substrate specificity, while APases involved in carbon metabolism (i.e. phosphoglycolate phosphatase, 3-PGA phosphatase and phosphoenolpyruvate (PEP) phosphatase) have much stricter substrate specificities (Duff *et al.*, 1991; del Pozo *et al.*, 1999; Miller *et al.*, 2001). Both intra- and extracellular APases are prominent in plants, and their activities have traditionally been used as markers for P-deficiency (Duff *et al.*, 1994; Baldwin *et al.*, 2001). Intracellular forms are found in the vacuole or cytoplasm. Although extracellular APases occur in the root apoplast and are frequently released from cell suspension cultures, extracellular types are usually localized in the cell wall, outer surface of root epidermal cells, and the root apical meristem. Intracellular APases appear to be much less stable than extracellular forms, which remain stable for hours to days (Goldstein *et al.*, 1988; Duff *et al.*, 1989a,

1991; Miller *et al.*, 2001). Increased stability would, of course, be a critical characteristic of any APase excreted into the rhizosphere. While intracellular APases induced during P deficiency function primarily in the release of P from senescent tissue for remobilization and in bypassing the P-requiring steps in C metabolism (Duff *et al.*, 1989b; Plaxton & Carswell, 1999), root extracellular APases are more likely to be involved in acquisition from soil (Marschner *et al.*, 1986; Tarafdar & Claassen, 1988). Although the acquisition of P from soil organic matter is equivocal, Miller *et al.* (2001) postulated that secretion of even low-affinity P-cleaving enzymes into soils with high organic P is an effective mechanism to provide additional sources of P for plant growth. Here, we focus on recent progress in understanding biochemical and molecular aspects of APases with special emphasis on exuded APases from cluster roots of P-deficient white lupin.

The literature abounds with reports of APases secreted from roots of P-deficient plants (Goldstein *et al.*, 1988; Li *et al.*, 1997a,b; Gilbert *et al.*, 1999; Hayes *et al.*, 1999; Zhang & McManus, 2000; Hunter & Leung, 2001). However, only recently have APase genes and proteins from *Arabidopsis*, tomato, and lupin cluster roots been fully characterized (del Pozo *et al.*, 1999; Wasaki *et al.*, 1999, 2002; Haran *et al.*, 2000; Baldwin *et al.*, 2001; Miller *et al.*, 2001). del Pozo *et al.* (1999) isolated and characterized a type 5 purple acid phosphatase, *AtACP5*, gene from *Arabidopsis*. The gene contained a 1014-bp open reading frame (ORF) encoding 338 amino acid residues. The protein had a 31 amino acid *N*-terminal extension with characteristics of a membrane targeting signal peptide. Both roots and shoots showed highly induced expression of *AtACP5* under P deficiency and addition of P could reverse induction. Induction of *AtACP5* also occurred in senescent tissues and in response to salt stress. Although *AtACP5* was highly expressed in roots under P-stress, the authors could find no evidence for secretion of *AtACP5* protein. They concluded that the protein was tightly anchored to the cell wall or plasmalemma. The *AtACP5* promoter, when fused to GUS and used as a reporter, showed high activity in response to P-deficiency, senescence, and salt stress. They concluded that *AtACP5* was involved more in internal P remobilization than P acquisition. Another APase (*LePS2*) implicated in internal remobilization of P has been characterized from tomato (Baldwin *et al.*, 2001). *LePS2* contains an 810-bp ORF encoding a 269 amino acid protein. The protein contains no 5'-signal peptide and is postulated to be in the cytosol. *LePS2* was highly induced in all tissues in P-deficient plants and expression was not induced by other nutrient stresses. Induction of *LePS2* transcripts could also be reversed by addition of P to stressed plants. Both *AtACP5* and *LePS2* were rapidly induced under P-deficient conditions, suggesting that plant molecular responses to P are tightly controlled.

The first truly secreted APase (sAPase) gene to be characterized was from *Arabidopsis* (*AtsAPase*; Haran *et al.* 2000).

The *AtsAPase* transcript is 1380 bp corresponding to a 46 kDa protein of approximately 450 amino acids. This protein contains a 34 amino acid *N*-terminal extension predicted to target the mature protein to the endomembrane system. The gene was highly induced in roots of P-deficient plants. When the *AtsAPase* transit sequence was fused to green fluorescent protein (GFP) as a reporter, GFP was detected in root exudates of P-deficient plants. The authors proposed that the APase transit sequence would be useful in strategies to produce and obtain relatively pure proteins of industrial importance from plant roots.

Root APase activity increases dramatically in P-deficient white lupin plants (Ozawa *et al.*, 1995; Gilbert *et al.*, 1999). This large increase in APase activity is associated with the appearance of a novel isoform (Wasaki *et al.*, 1997; Gilbert *et al.*, 1999). Moreover, this novel form induced by P-stress is also secreted into the rhizosphere of cluster roots (Gilbert *et al.*, 1999; Miller *et al.*, 2001; Wasaki *et al.*, 2002). When cluster roots from P-deficient plants are placed on agar containing APase substrate, exudation of APase activity into the rhizosphere can be detected within minutes (Fig. 7). White lupin sAPase is synthesized as a preprotein having an M_r of 52 kDa. During secretion the protein's 31-amino acid presequence is cleaved, giving rise to a 49 kDa processed protein. Similar to other APases, the sAPase is a glycoprotein. Protein blots probed with antibodies for the sAPase showed rapid accumulation of the protein in P-deficient roots accompanied by secretion into the rhizosphere (Miller *et al.*, 2001). The cDNA for white lupin sAPase contained a 1380 bp open reading frame capable of coding a 460-amino acid protein. Expression of sAPase mRNAs was quite specific (Fig. 2) and found predominantly under P-deficiency in cluster roots (Miller *et al.*, 2001; Wasaki *et al.*, 2002). Slight induction was also found in Al stressed roots. Miller *et al.* (2001) isolated the promoter for sAPase and showed that the gene was responsive to internal P concentration. In addition, the promoter was functional in alfalfa, showing that the molecular events controlling plant response to P deficiency are conserved between plants. The fact that a 50 bp region within the promoter of white lupin sAPase is 72% identical to a region in the *Arabidopsis* P-induced APase supports this interpretation. White lupin's ability to access P unavailable to other species (Braun & Helmke, 1995) may not only reside in its exudation of organic acids but may also be a consequence of exuded APase activity (Fig. 3b).

VIII. Genetic regulation of P responsive genes

As mentioned previously, numerous molecular events are set in motion in response to P deficiency resulting in biochemical changes that facilitate P conservation and/or acquisition. Raghothama (1999) has suggested that more than 100 genes may be involved in plant response to low-P stress. The fact that many of the molecular and biochemical changes in




Fig. 7 *In vivo* staining for acid phosphatase activity in roots of normal and cluster roots of white lupin. The root segments were embedded in an agarose–acid phosphatase substrate mixture and photographed at 15 min (a) and 25 h (b). The dark, purple color indicates acid phosphatase activity in the roots and the root exudates. Acid phosphatase is rapidly exuded into the rhizosphere of P-deficient plants; +, plants that are P sufficient; –, plants that are P deficient.

response to P deficiency occur in synchrony suggests that the genes involved are coordinately expressed and share a common regulatory system. Raghothama (1999) and others (Goldstein, 1992; Schachtman *et al.*, 1998) have suggested that regulation of plant response to P stress may reside in a phosphate (*PHO*) regulatory system similar to that found in yeast and bacteria. The *PHO* regulon in bacteria and yeast represents a complex multigene system having both structural and regulatory components (Vogel & Hinnen, 1990; Wanner, 1993; Lenburg & O'Shea, 1996; Oshima, 1997). The P_i signaling in these organisms involves a classic two-component system which includes a sensor protein and a response regulator protein. The sensor element is ascribed to a transmembrane protein comprised of an extracellular module, involved in sensing environmental cues, fused to a protein kinase module, capable of autophosphorylation. The response regulator is cytosolic and also has two modules, one of which has a phosphorylation site, and the other acts as a transcription (*trans*) factor. Induction of -P response genes, in simplest terms, involves the sensor protein receiving the environmental cue and catalysing autophosphorylation. The kinase module of the sensor then phosphorylates the receiver module of the response regulator. In turn, the *trans*-factor module of the response regulator is then rendered capable of binding to specific *cis*-elements in the promoters of P responsive genes. Those specific promoter *cis*-elements in P responsive genes that bind with the *PHO* response regulator *trans*-factor are called *PHO*-boxes.

In recent years, plant two-component signaling systems have been identified that are involved in perception of and response to ethylene, cytokinin and osmoticum (for reviews

see Sakakibara *et al.*, 1999; Urao *et al.*, 2000; Hwang & Sheen, 2001; Lohrmann & Harter, 2002). By contrast to the traditional prokaryotic two-component signaling system which comprises two proteins, the two-component system of plants appears to be more complex, involving a multistep process and at least three proteins (Sakakibara *et al.*, 2000; Urao *et al.*, 2000; Chang & Stadler, 2001; Gilroy & Trewavas, 2001; Lohrmann & Harter, 2002). The fundamental system in plants (Fig. 8) involves perception by the input domain of a membrane bound sensor histidine (His) kinase which auto-phosphorylates an aspartate (Asp) residue in the transmitter domain. The transmitter domain then phosphorylates a His residue on a phosphotransfer (HPT) protein which shuttles into the nucleus and transfers the P to an Asp residue of the response regulator (RR). Upon phosphorylation, the RR undergoes conformational change enabling it to bind to *cis*-elements within the promoter of genes to be activated by the signal molecule. More than 15 His-Asp kinases, five HPT, and 16 RR proteins have been identified in plants (Urao *et al.*, 2000; Lohrmann & Harter, 2002). Although this multistep two-component signaling mechanism has yet to be linked to signaling low and/or high P, it has been implicated in signaling N (Sakakibara *et al.*, 1999) and osmotic (Urao *et al.*, 1999) stress.

A growing body of evidence derived from studies of P-starvation response defective mutants of *Chlamydomonas* (Wykoff *et al.*, 1999) and *Arabidopsis* (Rubio *et al.* 2001) as well as soybean (*Glycine max*) P response genes (Tang *et al.*, 2001) indicates that nuclear localized transcription factors are involved in regulation of plant response to P. Genetic analysis of the *Chlamydomonas* and *Arabidopsis* P response mutants




Fig. 8 Postulated two-component signaling system for perceiving and responding to environmental stimuli. This multistep, signal-response system is thought to involve three proteins: (1) a histidine kinase input-receiver protein undergoes autophosphorylation in response to an environmental (i.e. low P, N, O₂) signal; (2) the phosphate (P) is then transferred to a phosphorelay protein which has a nuclear targeting signal; (3) upon entering the nucleus the P is transferred from the phosphorelay protein to the response-regulator (RR) protein which undergoes conformational change allowing it to bind to specific *cis*-elements within the promoter of downstream genes involved in the plant's response to the environment.

showed that the mutations had occurred in MYB coiled-coil transcription factors designated *PSR1* and *PHR1*, respectively. Evaluation of the *Arabidopsis* genome shows the presence of 15 MYB coiled-coil proteins. Although *PSR1* and *PHR1* transcripts are detected independently of the P status of the plant, both increase under P-deficiency. Both MYB genes are proposed to act downstream from the initial P signaling recognition event. The *Arabidopsis* *PHR1* MYB protein was shown to bind to an imperfect palindromic consensus sequence, 5'-GNATATNC-3', in the promoters of several P starvation-responsive genes including white lupin *sAPase*, *LaPT1*, and *LaMYB1* (Liu *et al.*, 2001; Miller *et al.*, 2001). By comparison, the soybean vegetative storage protein (VSPB) which is downregulated in response to added P was shown to be regulated in part by homeodomain leucine zipper protein (HD-ZIP) which binds to the CATTAATTAG element in the VSPB promoter.

Raghothama (1999) has noted that PHO-box like elements occur in some P-deficiency induced genes and has recently identified a *cis*-element 5'-CACGT(G/T)-3' in the promoter of several P-response genes. This element is identical to the Pho4 binding domain in yeast P-response genes (Ogawa *et al.*, 1995). We have also recently identified this motif in the promoters of P-deficiency induced APase (Miller

et al., 2001) and a MYB transcription factor (J-Q Liu & C. P. Vance unpubl. data), genes which are highly expressed in cluster roots. Interestingly, CACGTG is known as a G-box in plant genes which are regulated by various biotic and abiotic stresses. Also, relevant to identifying genes affecting nutrient acquisition and root development, Zhang and Forde (1998, 2000) have isolated and functionally characterized a MADs box transcription factor, *ANRI*, that controls nitrate-induced changes in lateral root development. Thus, there are likely many transcription factors and *cis*-elements that interact to control nutrient acquisition and root architecture. Many of these may be active during the development of white lupin cluster roots as evidenced by our sequencing of ESTs from these organs revealing *TUBBY*, *GRAB/NAM*, *RAS*, *DIMINUTO*, and *SCARECROW*-like genes (<http://www.home.earthlink.net/~whitelupinacelimation>).

IX. Improving P acquisition

Strategies for improving P acquisition and use involve both traditional plant breeding approaches and transgenic technology. Improvement through either approach is dependent upon: (1) identifying a suitable plant trait or gene that potentially affects acquisition and/or use; (2) finding or generating sufficient genetic variation for traits within a germplasm source; and (3) introgression of the trait into agronomically suitable backgrounds. Suitable plant traits having potential to impact P acquisition and use include: root architecture (Yan, 1999; Lynch & Brown, 2001; Williamson *et al.*, 2001), organic acid synthesis and extrusion (Hoffland *et al.*, 1992; Hinsinger & Gilkes, 1997; Neumann & Römheld, 1999), exudation of APases and phytases (Hayes *et al.*, 1999; Miller *et al.*, 2001; Richardson *et al.*, 2001; Brinch-Pedersen *et al.*, 2002); and phosphate transporters (Raghothama, 1999; Gilroy & Jones, 2000; Vance, 2001).

Roots have been notoriously recalcitrant to study and to improve through genetic selection due to the fact that their development is affected by a multitude of genes and their subterranean nature. However, as outlined here and elsewhere in this review, advances in understanding the genetic control of roots, improved methods for root evaluation, and reaffirmation of their importance for improving nutrient use efficiency have led to a renaissance of discovery in root biology that should result in significant improvements to crop plants.

Genetic variability has been documented in a wide range of species (and probably exists in all species) for tap root elongation, basal root growth angle (i.e. rooting depth, lateral and adventitious root development, and root hair numbers; Klepper, 1992; Charlton, 1996). Inclusively these traits comprise root architecture and each has been identified as important to P acquisition (Lynch, 1995; Eshel *et al.*, 1996; Gahoonia & Nielsen, 1997; Care & Caradus, 1999; Yan, 1999; Narang *et al.*, 2000). *Phaseolus vulgaris* has been the prototype for selection of root architecture ideotypes that result in improved

P acquisition (Lynch, 1995; Yan, 1999; Lynch & Brown, 2001). Genetic selection as well as root simulation models with bean have shown enhanced P uptake in genotypes with shallow roots, accompanied by heightened numbers of lateral roots with abundant root hairs. Selection for increased root hair abundance has led to enhanced P acquisition in bean, clover, barley and *Arabidopsis*. Wissuwa & Ae (2001) have identified four quantitative trait loci (QTL) in rice (*Oryza sativa*) associated with increased P uptake under P-deficiency, two of which were related to maintenance of root growth under low P. Similarly, these root architecture traits are also important in the acquisition of N and other nutrients (Klepper, 1992; Zhang & Forde, 1998; Lamb *et al.*, 2000; Vance & Lamb, 2001).

Phosphorus acquisition has also been improved through approaches aimed at increasing organic acid synthesis in and/or exudation from plant roots. This approach is based upon the large body of evidence showing that exudation of citrate and malate from roots effectively solubilizes unavailable sources of P (Marschner *et al.*, 1986). Release of organic acids into the rhizosphere in exudates leads to metal chelation and subsequent desorption of P_i from the soil matrix, with a concomitant increase in availability. Rhizosphere acidification by protons and/or organic acids may be an accompanying trait when root architecture is modified through selection as shown in bean (Yan, 1999), *Arabidopsis* (Narang *et al.*, 2000), rice (Yang *et al.*, 2000) and pigeon pea (Ae *et al.*, 1990). The acidification of the rhizosphere of white lupin cluster roots also exemplifies the concurrent exudation of organic acids accompanying a change in root development. Overexpression of genes involved in organic acid synthesis appears to be a useful strategy to stimulate P acquisition by enhancing exudation of citrate and malate. Koyama *et al.* (1999) developed carrot (*Daucus carota*) cell lines that overexpressed mitochondrial citrate synthase. The overexpressing cell lines exuded more citrate and had improved growth on $AlPO_4^-$ medium compared with untransformed controls. In a similar approach but using a bacterial citrate synthase gene driven by the CaMV^{35S} promoter, Lopez-Bucio *et al.* (2000a) demonstrated that overexpression of citrate synthase resulted in increased secretion of citrate into the rhizosphere and enhanced P accumulation. Our laboratory recently overexpressed a novel form of MDH in alfalfa (Tesfaye *et al.*, 2001). Transgenic plants containing the novel MDH driven by the CaMV^{35S} promoter had increased organic acid synthesis and exudation, which resulted in increased P accumulation compared with either transformed or untransformed controls. Interestingly, in each of the above cases where overexpression of genes involved in citrate and malate production enhanced organic acid synthesis and P acquisition, the transformed plants had greater Al^{3+} tolerance than controls (Lopez-Bucio *et al.*, 2000b; Tesfaye *et al.*, 2001). It is well established that Al^{3+} tolerance in plants is related to organic acid accumulation and/or exudation by roots (Kochian, 1995; Ryan *et al.*, 2001). Moreover, increased

organic acid synthesis and exudation has been implicated in tolerance to Pb and other metals (Mathys, 1977; Godbold *et al.*, 1984; Kochian, 1995; Yang *et al.*, 2000). In addition, accumulation and exudation of citrate and malate by roots has been implicated in alleviating Fe deficiency (de Vos *et al.*, 1986; Jones, 1998). Thus, genetic engineering of plants for enhanced organic acid synthesis may have far-reaching effects on plant nutrition and stress tolerance.

Other targets which have potential to improve P_i acquisition include high-affinity P_i transporters (Mitsukawa *et al.*, 1997) and root exuded phytase (Richardson *et al.*, 2001). Overexpression of the *Arabidopsis* high-affinity P_i transporter gene (*PHT1*) in tobacco resulted in cell lines with enhanced growth under P_i -limited conditions (Mitsukawa *et al.*, 1997). Transgenic tobacco lines overexpressing *PHT1* had threefold greater P_i uptake than controls and 50% greater growth under low P_i conditions. Richardson *et al.* (2001) generated transgenic *Arabidopsis* plants that expressed an *Aspergillus* phytase gene and exuded phytase into the rooting medium. The transgenic plants could grow on phytate as the sole source of P whereas control plants could not. We have recently developed alfalfa plants that can overexpress several diverse proteins in roots in efforts to metabolically engineer improved nutrient use acquisition (APase, MDH and metallothioneins). These transgenic plants are currently awaiting characterization.

Evidence is accumulating that selection for and overexpression of selected genes involved in either root development or P acquisition can improve nutrient acquisition. As the arsenal of plant genes involved in these processes expands through whole-genome approaches, we will undoubtedly see more successful examples of plants with better nutrient acquisition and utilization created through traditional breeding and transgenic technology. Because of the urgent need for plant germplasm with improved P-use efficiency, it is imperative that research programs mount multidisciplinary teams that combine traditional plant breeding, biotechnology and plant physiology.

X. Synopsis

The world is on the brink of a new agriculture, one that involves the marriage of plant biology and agroecology under the umbrella of biotechnology and germplasm improvement. Although P fertilizers will continue to play a major role in intensive agriculture, depletion of natural resources, loss of biodiversity, and long-term unsustainability require that alternative strategies be investigated and implemented to buffer against food insecurity and environmental degradation. Furthermore, because improved P acquisition and use by plants has immediate and direct benefit in extensive agriculture in developing countries where access to fertilizers is limited, funding for research at international centers should be a high priority. The following recommendations deserve attention: (1) expand research in root biology;

(2) continue to isolate, characterize and develop a fundamental understanding of individual genes holding promise for application to improving P nutrition; (3) enhance the expression of genes and increase the synthesis of gene products, such as those involved in transport of nutrients and exudation of organic acids, through transgenic technology and incorporate these traits into adapted germplasm; (4) continue and expand collaborations between plant breeders and physiologists to develop germplasm with enhanced P-use efficiency; and (5) reemphasize applied research programs directed toward sustainable approaches to enhancing soil P through crop management (such as intercropping, rotations and incorporation of legumes).

Acknowledgements

This work was funded in part by the United States Department of Agriculture, National Research Initiative Grant No. USDA-CSREES/98-35100-6098 and USDA-CSREES/2002-35100-12206, and United States Department of Agriculture, Agricultural Research Service CRIS no. 3640-21000-019-00D.

References

- Abelson PH. 1999. A potential phosphate crisis. *Science* 283: 2015.
- Adams MA, Pate JS. 2002. Phosphorus sources and availability modify growth and distribution of root clusters and nodules of native Australian legumes. *Plant, Cell & Environment* 26: 837–850.
- Ae N, Arihara J, Okada K, Yoshihara T, Johansen C. 1990. Phosphorus uptake by pigeon pea and its role in cropping systems of the Indian subcontinent. *Science* 248: 477–480.
- Aono T, Kanada N, Ijima A, Oyaizu H. 2001. The response of the phosphate uptake system and the organic acid exudation system to phosphate starvation in *Sesbania rostrata*. *Plant and Cell Physiology* 42: 1253–1264.
- Ashihara H, Li X-N, Ukaji T. 1988. Effect of inorganic phosphate on the biosynthesis of purine and pyrimidine nucleotides in suspension-cultured cells of *Catharanthus roseus*. *Annals of Botany* 61: 225–232.
- Avio L, Sbrana C, Giovannetti M. 1990. The response of different species of *Lupinus* to VAM endophytes. *Symbiosis* 9: 321–323.
- Baldwin JC, Athikkattuvalasu SK, Raghothama KG. 2001. LEPS2, a phosphorus starvation-induced novel acid phosphatase from tomato. *Plant Physiology* 125: 728–737.
- Barber SA, Walker JM, Vasey EH. 1963. Mechanisms for the movement of plant nutrients from the soil and fertilizer to the plant root. *Journal of Agricultural and Food Chemistry* 11: 204–207.
- Barbier-Brygoo H, Vinauger M, Colcombet J, Ephritikhine G, Frachisse J, Maurel C. 2000. Anion channels in higher plants: functional characterization, molecular structure and physiological role. *Biochimica et Biophysica Acta* 1465: 199–218.
- Baskin TJ, Cork A, Williamson RE, Gorst JR. 1995. *STUNTEDPLANT1*, a gene required for expansion in rapidly elongating but not dividing cells mediating root growth responses to applied cytokinin. *Plant Physiology* 107: 233–243.
- Bates TR, Lynch JP. 2000. The efficiency of *Arabidopsis thaliana* (Brassicaceae) root hairs in phosphorus acquisition. *American Journal of Botany* 87: 964–970.
- Benning C. 1998. Biosynthesis and function of the sulfolipid sulfoquinovosyl diacylglycerol. *Annual Reviews of Plant Physiology and Plant Molecular Biology* 49: 53–75.
- Bielecki RL. 1973. Phosphate pools, phosphate transport, and phosphate availability. *Annual Review of Plant Physiology* 24: 225–252.
- Binns AN. 1994. Cytokinin accumulation and action: biochemical, genetic, and molecular approaches. *Annual Review of Plant Physiology and Plant Molecular Biology* 45: 173–196.
- Biswas TK, Cundiff C. 1991. Multiple forms of acid phosphatase in germinating seeds of *Vigna sinensis*. *Phytochemistry* 30: 2119–2125.
- Blakely LM, Blakely RM, Colowit PM, Elliot DS. 1988. Experimental studies on lateral root formation in radish seedling roots. II. Analysis of the dose–response to exogenous auxin. *Plant Physiology* 87: 414–419.
- Boerjan W, den Boer B, van Montagu M. 1992. Molecular genetic approaches to plant development. *International Journal of Developmental Biology* 36: 59–66.
- Boerjan W, Cervera MT, Delarue M, Beeckman T, Dewitte W, Bellini C, Caboche M, van Onckelen H, van Montagu M, Inze D. 1995. Superroot, a recessive mutation in *Arabidopsis*, confers auxin overproduction. *Plant Cell* 7: 1405–1419.
- Borch K, Bouma TJ, Lynch JP, Brown KM. 1999. Ethylene: a regulator of root architectural responses to soil phosphorus availability. *Plant, Cell & Environment* 22: 425–431.
- Braum SM, Helmke PA. 1995. White lupin utilizes soil phosphorus that is unavailable to soybean. *Plant and Soil* 176: 95–100.
- Brinch-Pedersen H, Sorenson LD, Holm PB. 2002. Engineering crop plants: getting a handle on phosphate. *Trends in Plant Science* 7: 118–125.
- Brown MH, Paulsen IT, Skurray RA. 1999. The multidrug efflux protein NorM is a prototype of a new family of transporters. *Molecular Microbiology* 31: 394–395.
- Bryce JH, Azcon-Bieto J, Wiskich JT, Day DA. 1990. Adenylate control of respiration in plants: the contribution of rotenone insensitive electron transport to ADP-limited oxygen consumption by soybean mitochondria. *Physiologia Plantarum* 78: 105–111.
- Bumb BL, Baanante CA. 1996. *The role of fertilizer in sustaining food security and protecting the environment. Food, agriculture and the environment discussion paper 17*. Washington, DC, USA: International Food Policy Research Institute.
- Burleigh SH, Cavagnaro T, Jakobsen I. 2002. Functional diversity of arbuscular mycorrhizas extends to the expression of plant genes involved in P nutrition. *Journal of Experimental Botany* 53: 1593–1601.
- Cao Y, Ward JM, Kelly WB, Ichida AM, Gaber RF, Anderson JA, Uozumi N, Schroeder JI, Crawford NM. 1995. Multiple genes, tissue specificity, and expression-dependent modulation contribute to the functional diversity of potassium channels in *Arabidopsis thaliana*. *Plant Physiology* 109: 1093–1106.
- Care DA, Caradus JR. 1999. Root hairs of white clover genotypes differing in phosphorus response. In: Lynch JP, Deikman J, eds. *Phosphorus in plant biology: regulatory roles in molecular, cellular, organismic, and ecosystem processes*. Rockville, MD, USA: American Society of Plant Physiology, 316–317.
- Cary AJ, Liu W, Howell SH. 1995. Cytokinin action is coupled to ethylene in its effect on the inhibition of root and hypocotyl elongation in *Arabidopsis thaliana* seedlings. *Plant Physiology* 107: 1075–1082.
- Cary AJ, Uttamchandani SJ, Smets R, van Onckelen HA, Howell SH. 2001. *Arabidopsis* mutants with increased organ regeneration in tissue culture are more competent to respond to hormonal signals. *Planta* 213: 700–707.
- Casimiro I, Marchant A, Bhalerao RP, Beeckman T, Dhooge S, Swarup R, Graham N, Inze D, Sandberg G, Casero PJ, Bennett MJ. 2001. Auxin transport promotes *Arabidopsis* lateral root initiation. *Plant Cell* 13: 843–852.
- Celenza JL Jr, Grisafi PL, Fink GR. 1995. A pathway for lateral root formation in *Arabidopsis thaliana*. *Genes and Development* 9: 2131–2142.
- Chang C, Stadler R. 2001. Ethylene hormone receptor action in *Arabidopsis*. *Bioessays* 23: 619–627.

- Charlton WA. 1996. Lateral root initiation. In: Waisel Y Eshel A Kafkafa U, eds. *Plant roots: the hidden half*, 2nd edn. New York, NY, USA: Marcel Dekker. 149–173.
- Chiou TJ, Liu H, Harnson MJ. 2001. The spatial expression patterns of a phosphate transporter (MtPT1) from *Medicago truncatula* indicate a role in phosphate transport at the root/soil interface. *Plant Journal* 25: 281–293.
- Cho H-T, Cosgrove DJ. 2002. Regulation of root hair initiation and expansin gene expression in Arabidopsis. *Plant Cell* 14: 3237–3253.
- Christeller JT, Tolbert NE. 1978. Phosphoglycolate phosphatase: purification and properties. *Journal of Biological Chemistry* 253: 1780–1785.
- Christianson ML, Warnick DA. 1985. temporal requirements for phytohormone balance in the control of organogenesis *in vitro*. *Developmental Biology* 112: 494–497.
- Comerford NB. 1998. Soil phosphorus bioavailability. In: Lynch JP Deikman J, eds. *Phosphorus in plant biology: regulatory roles in molecular, cellular, organismic, and ecosystem processes*. Rockville, MD, USA: American Society of Plant Physiology, 136–147.
- Council for Agricultural Science and Technology (CAST). 1988. *Long term viability of US agriculture. Report no. 114*. Ames, IA, USA: CAST.
- Curl EA, Truelove B. 1986. *The Rhizosphere*. Berlin, Germany: Springer-Verlag.
- Dancer J, Veith R, Feil R, Komor E, Stitt M. 1990. Independent changes of inorganic pyrophosphate and the ATP/ADP or UTP/UDP ratios in plant suspension cultures. *Plant Science* 66: 59–63.
- Daram P, Brunner S, Persson BL, Amrhein N, Bucher M. 1998. Functional analysis and cell-specific expression of a phosphate transporter from tomato. *Planta* 206: 225–233.
- Day DA, Wiskich JT. 1995. Regulation of alternative oxidase activity in higher plants. *Journal of Bioenergetics and Biomembranes* 27: 379–385.
- Debeaujon I, Peeters AJ, Leon-Kloosterziel KM, Koornneef M. 2001. The TRANSPARENT TESTA12 gene of *Arabidopsis* encodes a multidrug secondary transporter-like protein required for flavonoid sequestration in vacuoles of the seed coat endothelium. *Plant Cell* 13: 853–871.
- Delhaize E, Ryan PR. 1995. Aluminum toxicity and tolerance in plants. *Plant Physiology* 107: 315–321.
- Delhaize E, Ryan PR, Randall PJ. 1993. Aluminum tolerance in wheat (*Triticum aestivum* L.). II. Aluminum-stimulated excretion of malic acid from root apices. *Plant Physiology* 103: 695–702.
- Diener AC, Gaxiola RA, Fink GR. 2001. Arabidopsis ALF5, a multidrug efflux transporter gene family member, confers resistance to toxins. *Plant Cell* 13: 1625–1638.
- Dinkelaker B, Hengeler C, Marschner H. 1995. Distribution and function of proteoid roots and other root clusters. *Acta Botanica* 108: 169–276.
- Dinkelaker B, Romheld V, Marschner H. 1989. Citric acid excretion and precipitation of calcium citrate in the rhizosphere of white lupin (*Lupinus albus* L.). *Plant, Cell & Environment* 12: 285–292.
- Dolan L. 2001. The role of ethylene in root hair growth in Arabidopsis. *Journal of Plant Nutrition and Soil Science* 164: 141–145.
- Drew MC. 1975. Comparison of the effect of a localized supply of phosphate, nitrate, ammonium and potassium on the growth of the seminal root system, and the shoot, in barley. *New Phytologist* 75: 479–490.
- Duff SMG, Lefebvre DD, Plaxton WC. 1989a. Purification and characterization of a phosphoenolpyruvate phosphatase from *Brassica nigra* suspension cells. *Plant Physiology* 90: 734–741.
- Duff SMG, Moorhead GBG, Lefebvre DD, Plaxton WC. 1989b. Phosphate starvation inducible 'bypasses' of adenylate and phosphate dependent glycolytic enzymes in *Brassica nigra* suspension cells. *Plant Physiology* 90: 1275–1278.
- Duff SM, Plaxton WC, Lefebvre DD. 1991. Phosphate-starvation response in plant cells: de novo synthesis and degradation of acid phosphatases. *Proceedings of the National Academy of Sciences, USA* 88: 9538–9542.
- Duff SMG, Sarath G, Plaxton WC. 1994. The role of acid phosphatases in plant phosphorus metabolism. *Physiologia Plantarum* 90: 791–800.
- Dumont F, Joris B, Gumusboga A, Bruyninx M, Loppes R. 1993. Isolation and characterization of cDNA sequences controlled by inorganic phosphate in *Chlamydomonas reinhardtii*. *Plant Science* 89: 55–67.
- Ellington CP. 1999. Crop yield responses to phosphorus. In: Ellington CP, ed. *Phosphorus for agriculture: a situation analysis*. Norcross, GA, USA: Potash/Phosphate Institute, 25–41.
- Epstein E. 1953. Mechanism of ion absorption by roots. *Nature* 171: 83–84.
- Eshel A, Nielsen KL, Lynch JP. 1996. Screening bean (*Phaseolus vulgaris* L.) genotypes for differences in phosphorus utilization efficiency. *Israel Journal of Plant Science* 44: 225–233.
- Essigmann B, Güler S, Narang RA, Linke D, Benning C. 1998. Phosphate availability affects the thylakoid lipid composition and the expression of *SQD1*, a gene required for sulfolipid biosynthesis in *Arabidopsis thaliana*. *National Academy of Sciences, USA* 95: 1950–1955.
- Faure J-D, Vittorioso P, Santoni V, Fraiser V, Prinsen E, Barlier I, van Onckelen HA, Caboche M, Bellini C. 1998. The *PASTICCINO* genes of *Arabidopsis thaliana* are involved in the control of cell division and differentiation. *Development* 125: 909–918.
- Fukaki H, Tameda S, Masuda H, Taska M. 2002. Lateral root formation is blocked by a gain-of-function mutation in the SOLITARY-ROOT/IAA14 gene of *Arabidopsis*. *Plant Journal* 29: 153–168.
- Furihata T, Suzuki M, Sakurai H. 1992. Kinetic characterization of two phosphate uptake systems with different affinities in suspension-cultured *Catharanthus roseus* protoplasts. *Plant and Cell Physiology* 33: 1151–1157.
- Gahoonia TS, Nielsen NE. 1997. Variation in root hairs of barley cultivars doubled soil phosphorus uptake. *Euphytica* 98: 177–182.
- Gardner W, Parbery D, Barber D. 1981. Proteoid root morphology and function in *Lupinus albus*. *Plant and Soil* 60: 143–147.
- Gardner WK, Parbery DG, Barber DA. 1982. The acquisition of phosphorus by *Lupinus albus* L. I. Some characteristics of the soil/root interface. *Plant and Soil* 68: 19–32.
- Gardner WK, Barber DA, Parbery DG. 1983. The acquisition of phosphorus by *Lupinus albus* L. III. The probable mechanism by which phosphorus movement in the soil/root interface is enhanced. *Plant and Soil* 70: 107–124.
- Ge Z, Rubio G, Lynch JP. 2000. The importance of root gravitropism for inter-root competition and phosphorus acquisition efficiency: results from a geometric simulation model. *Plant and Soil* 218: 159–171.
- Geelen D, Lurin C, Bouchez D, Frachisse JM, Lelievre F, Courtial B, Barbier-Brygoo H, Maurel C. 2000. Disruption of putative anion channel gene AtCLC-a in Arabidopsis suggests a role in the regulation of nitrate content. *Plant Journal* 21: 259–267.
- Gerke J. 1994. Kinetics of soil phosphate desorption as affected by citric acid. *Zeitschrift für Pflanzenernährung und Bodenkunde* 157: 17–22.
- Gerke JR, Römer W, Jungk A. 1994. The excretion of citric and malic acid by proteoid roots of *Lupinus albus* L. effect on soil solution concentrations of phosphate, iron, and aluminum in the proteoid rhizosphere in samples of an oxisol and a lurisol. *Zeitschrift für Pflanzenernährung und Bodenkunde* 157: 289–294.
- Gilbert GA, Knight CD, Vance CP, Allan DL. 1997. Does auxin play a role in the adaptations of white lupin roots to phosphate deficiency. *To Plant Physiology* 114: 31.
- Gilbert GA, Knight JD, Vance CP, Allan DL. 1999. Acid phosphatase activity in phosphorus-deficient white lupin roots. *Plant, Cell & Environment* 22: 801–810.
- Gilbert GA, Knight JD, Vance CP, Allan DL. 2000. Proteoid root development of phosphorus deficient lupin is mimicked by auxin and phosphonate. *Annals of Botany* 85: 921–928.
- Gilbert GA, Vance CP, Allan DL. 1998. Regulation of white lupin metabolism by phosphorus availability. In: Lynch JP Deikman J, eds. *Phosphorus in plant biology: regulatory roles in molecular, cellular, organismic and ecosystem processes*. Rockville, MD, USA: American Society of Plant Physiology, 157–167.
- Gilroy S, Jones DL. 2000. Through form to function: root hair development and nutrient uptake. *Trends in Plant Science* 5: 56–60.

- Gilroy S, Trewavas A. 2001. Signal processing and transduction in plant cells: the end of the beginning. *Nature Reviews Molecular and Cell Biology* 2: 307–314.
- Godbold DL, Horst WJ, Collins JC, Thurmman DA, Marschner H. 1984. Accumulation of zinc and organic acids in roots of zinc tolerant and non-tolerant ecotypes of *Deschampsia caespitosa*. *Journal of Plant Physiology* 116: 59–69.
- Goldstein AH. 1992. Phosphate starvation inducible enzymes and proteins in higher plants. In: Wray JL, ed. *Society for Experimental Biology Series 49: inducible plant proteins*. Cambridge, UK: Cambridge University Press, 25–44.
- Goldstein AH, Baertlein DA, McDaniel RG. 1988. Phosphate starvation inducible metabolism in *Lycopersicon esculentum*. I. Excretion of acid phosphatase by tomato plants and suspension-cultured cells. *Plant Physiology* 87: 711–715.
- Grierson P. 1992. Organic acids in the rhizosphere of *Banksia integrifolia* L. f. *Plant and Soil* 144: 259–265.
- Grierson P, Attwill P. 1989. Chemical characteristics of the proteoid root mat of *Banksia integrifolia* L. *Australian Journal of Botany* 37: 137–143.
- Grierson CS, Parker JS, Kemp AC. 2001. *Arabidopsis* genes with roles in root hair development. *Journal of Plant Nutrition and Soil Science* 164: 131–140.
- Haran S, Logendra S, Saskar M, Bratanova M, Raskin I. 2000. Characterization of *Arabidopsis* acid phosphatase promoter and regulation of acid phosphatase expression. *Plant Physiology* 124: 615–626.
- Harrison MJ. 1997. The arbuscular mycorrhizal symbiosis: an underground association. *Trends in Plant Science* 2: 54–60.
- Hayes JE, Richardson AE, Simpson RJ. 1999. Phytase and acid phosphatase activities in roots of temperate pasture grasses and legumes. *Australian Journal of Plant Physiology* 26: 801–809.
- Hechenberger M, Schwappach B, Fischer WN, Frommer WB, Jentsch TJ, Steinmeyer K. 1996. A family of putative chloride channels from *Arabidopsis* and functional complementation of a yeast strain with a CLC gene disruption. *Journal of Biological Chemistry* 271: 33632–33638.
- Hinsinger P. 2001. Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review. *Plant Soil* 237: 173–195.
- Hinsinger P, Gilkes RJ. 1997. Dissolution of rock phosphate in the rhizosphere of five plant species grown in an acid, P-fixing mineral substrate. *Geoderma* 75: 231–249.
- Hobbie L, Estelle M. 1995. The *axr4* auxin-resistant mutants of *Arabidopsis thaliana* define a gene important for root gravitropism and lateral root initiation. *Plant Journal* 7: 211–220.
- Hoefnagel MH, van Iren F, Libbenga KR. 1993. In suspension cultures of *Catharanthus roseus* the cyanide-resistant pathway is engaged in respiration by excess sugar in combination with phosphate or nitrogen starvation. *Physiologia Plantarum* 87: 297–304.
- Hoefnagel MH, Millar AH, Wiskich JT, Day DA. 1995. Cytochrome and alternative respiratory pathways compete for electrons in the presence of pyruvate in soybean mitochondria. *Archives of Biochemistry and Biophysics* 318: 394–400.
- Hoffland E, van den Boogaard R, Nelemans J, Findenegg G. 1992. Biosynthesis and root exudation of citric and malic acids in phosphate-starved rape plants. *New Phytologist* 122: 675–680.
- Holford ICR. 1997. Soil phosphorus: its measurement and its uptake by plants. *Australian Journal of Soil Research* 35: 227–239.
- Horgan JM, Wareing PF. 1980. Cytokinins and the growth responses of seedlings of *Betula pendula* Roth. & *Acer pseudoplatanus* L. to nitrogen and phosphorus deficiency. *Journal of Experimental Botany* 31: 525–532.
- Horst WJ, Kamh M, Jibrin JM, Chude VA. 2001. Agronomic measures for increasing P availability to crops. *Plant and Soil* 237: 211–233.
- Howitt SM, Udvardi MK. 2000. Structure, function and regulation of ammonium transporters in plants. *Biochimica et Biophysica Acta* 1465: 152–170.
- Hunter DA, Leung DW. 2001. Enhancement of acid phosphatase activity in *Capiscum annum* L. plants in response to phosphate deficiency. *Journal of Plant Nutrition and Soil Science* 164: 219–233.
- Hwang I, Sheen J. 2001. Two-component circuitry in *Arabidopsis* cytokinin signal transduction. *Nature* 413: 383–389.
- Jackson RB, Manwaring JH, Caldwell MM. 1990. Rapid physiological adjustment of roots to localized soil enrichment. *Nature* 344: 58–60.
- Jeschke W, Pate J. 1995. Mineral nutrition and transport in xylem and phloem of *Banksia prionotes* (Proteacea), a tree with dimorphic root morphology. *Journal of Experimental Botany* 46: 895–905.
- Johnson JF, Allan DL, Vance CP. 1994. Phosphorus stress-induced proteoid roots show altered metabolism in *Lupinus albus*. *Plant Physiology* 104: 657–665.
- Johnson JF, Allan DL, Vance CP, Weiblen G. 1996a. Root carbon dioxide fixation by phosphorus-deficient *Lupinus albus*. Contribution to organic acid exudation by proteoid roots. *Plant Physiology* 112: 19–30.
- Johnson JF, Vance CP, Allan DL. 1996b. Phosphorus deficiency in *Lupinus albus*. Altered lateral root development and enhanced expression of phosphoenolpyruvate carboxylase. *Plant Physiology* 112: 31–41.
- Johnston AM. 2002. Nutrient removal as a soil fertility planning tool. Fertilization Facts. <http://www.ppi-ppic.org>.
- Jones DL. 1998. Organic acids in the rhizosphere: a critical review. *Plant and Soil* 205: 25–44.
- Jungk A. 2001. Root hairs and acquisition of plant nutrients from soil. *Journal of Plant Nutrition and Soil Science* 164: 121–129.
- Jungk A, Asher CJ, Edwards DG, Meyer D. 1990. Influence of phosphate status on phosphate uptake kinetics of maize (*Zea mays*) and soybean (*Glycine max*). *Plant and Soil* 124: 175–182.
- Keerthisinghe G, Hocking PJ, Ryan PR, Delhaize E. 1998. Effect of phosphorus supply on the formation and function of proteoid roots of white lupin. *Plant, Cell & Environment* 21: 467–478.
- Kieber JJ. 1997. The ethylene response pathway in *Arabidopsis*. *Annual Review of Plant Physiology and Plant Molecular Biology* 48: 277–296.
- King JJ, Stimart DP, Fisher RH, Bleeker AB. 1995. A mutation altering auxin homeostasis and plant morphology in *Arabidopsis*. *Plant Cell* 7: 2023–2037.
- Klepper B. 1992. Development and growth of crop root systems. *Advances in Soil Science* 19: 1–25.
- Kochian LV. 1995. Cellular mechanisms of aluminum toxicity and resistance in plants. *Annual Review of Plant Physiology and Plant Molecular Biology* 46: 237–260.
- Koide RT, Kabir Z. 2000. Extraradical hyphae of the mycorrhizal fungus *Glomus intraradices* can hydrolyse organic phosphate. *New Phytologist* 148: 511–517.
- Kollmeier M, Dietrich P, Bauer CS, Horst WJ, Hedrich R. 2001. Aluminum activates a citrate-permeable anion channel in the aluminum-sensitive zone of the maize root apex. A comparison between an aluminum-sensitive and an aluminum-resistant cultivar. *Plant Physiology* 126: 397–410.
- Koyama H, Takita E, Kawamura A, Hara T, Shibata D. 1999. Over expression of mitochondrial citrate synthase gene improves the growth of carrot cells in Al-phosphate medium. *Plant and Cell Physiology* 40: 482–488.
- Kuiper D, Schuit J, Kuiper PJC. 1988. Effect of internal and external cytokinin concentrations on root growth and shoot to root ratio of *Plantago major* ssp. *pleiosperma* at different nutrient concentrations. *Plant and Soil* 111: 231–236.
- Lajtha K, Harrison AF. 1995. Strategies of phosphorus acquisition and conservation by plant species and communities. In: Tiessen H, ed. *Phosphorus in the global environment*. Chichester, UK: John Wiley Sons Ltd, 140–147.
- Lamb JFS, Samac DA, Barnes DK, Henjum KI. 2000. Increased herbage yield in alfalfa associated with selection for fibrous and lateral roots. *Crop Science* 40: 693–699.
- Lambers H. 1985. Respiration in intact plants and tissues: Its regulation and dependence on environmental factors, metabolism and invaded

- organisms. In: Douce R, Day DA, eds. *Encyclopedia of plant physiology, new series, higher plant cell respiration*, Vol. 18. Berlin, Germany: Springer Verlag, 418–473.
- Lambers H, Juniper D, Cawthray G, Veneklaas E, Martinez-Ferri E. 2002. The pattern of carboxylate exudation in *Banksia grandis* (Proteaceae) is affected by the form of phosphate added to the soil. *Plant and Soil* 238: 111–122.
- Lamont BB. 1973. Factors affecting the distribution of proteoid roots within the root system of two *Hakea* species. *Australian Journal of Botany* 20: 155–174.
- Lamont BB. 1982. Mechanisms for enhancing nutrient uptake in plants, with particular reference to Mediterranean South Africa and Australia. *Botanical Reviews* 48: 597–689.
- Lamont BB. 1993. Why are hairy root clusters so abundant in the most nutrient-impooverished soils of Australia? *Plant and Soil* 155/156: 269–272.
- Lamont BB, Brown G, Mitchel DT. 1984. Structure and environmental effects on the formation and function of proteoid roots in *Leucadendron laueolium*. *New Phytologist* 97: 381–390.
- Lance C, Cheauveau M, Dinzengremel P. 1985. The cyanide resistant pathway of plant mitochondria. In: Douce R Day DA, eds. *Encyclopedia of plant physiology, new series, higher plant cell respiration*, Vol. 18. Berlin, Germany: Springer Verlag, 202–247.
- Larsen PB, Degenhardt J, Tai CY, Stenzler LM, Howell SH, Kochian LV. 1998. Aluminum-resistant *Arabidopsis* mutants that exhibit altered patterns of aluminum accumulation and organic acid release from roots. *Plant Physiology* 117: 9–18.
- Lefebvre DD, Duff SMG, Fife CA, Julien-Inalsingh C, Plaxton WC. 1990. Response to phosphate deprivation in *Brassica nigra* suspension cells: enhancement of intracellular, cell surface and secreted acid phosphatase activities compared to increases in P_i -absorption rate. *Physiologia Plantarum* 93: 504–511.
- Leggiew G, Willmitzer L, Riesmeier JW. 1997. Two cDNAs from potato are able to complement a phosphate uptake-deficient yeast mutant: identification of phosphate transporters from higher plants. *Plant Cell* 9: 381–392.
- Lenburg ME, O'Shea EK. 1996. Signaling phosphate starvation. *Trends in Biochemical Sciences* 21: 383–387.
- Li M, Osaki M, Rao IM, Tadano T. 1997a. Secretion of phytase from roots of several species under phosphorus-deficient conditions. *Plant and Soil* 195: 161–169.
- Li M, Shinano T, Tadano T. 1997b. Distribution of exudates of lupin roots in the rhizosphere under phosphorus deficient conditions. *Soil Science and Plant Nutrition* 43: 237–245.
- Liao H, Rubio G, Yan X, Cao A, Brown KM, Lynch JP. 2001. Effect of phosphorus availability on basal root shallowness in common bean. *Plant and Soil* 232: 69–79.
- Liu C, Muchhal VS, Mukatira U, Kononowicz AK, Ragothama KG. 1998. Tomato phosphate transporter genes are differentially regulated in plant tissues by phosphorus. *Plant Physiology* 116: 91–99.
- Liu H, Trieu AT, Blaylock LA, Harrison MJ. 1998. Cloning and characterization of two phosphate transporters from *Medicago truncatula* roots: regulation in response to phosphate and to colonization by arbuscular mycorrhizal (AM) fungi. *Molecular Plant–Microbe Interactions* 11: 14–22.
- Liu J, Uhde-Stone C, Li A, Vance CP, Allan DL. 2001. A phosphate transporter with enhanced expression in proteoid roots of white lupin (*Lupinus albus* L.). *Plant and Soil* 237: 257–266.
- Lohrmann J, Harter K. 2002. Plant two-component signaling systems and the role of response regulators. *Plant Physiology* 128: 363–369.
- Lopez-Bucio J, de la Vega OM, Guevara-Garcia A, Herrera-Estrella L. 2000a. Enhanced phosphorus uptake in transgenic tobacco plants that overproduce citrate. *Nature Biotechnology* 18: 450–453.
- Lopez-Bucio J, Nieto-Jacobo MF, Ramirez-Rodriguez VV, Herrera-Estrella L. 2000b. Organic acid metabolism in plants: from adaptive physiology to transgenic varieties for cultivation in extreme soils. *Plant Science* 160: 1–13.
- Louis I, Racette S, Torrey JG. 1990. Occurrence of cluster roots on *Myrica cerifera* L. (Myricaceae) in water culture in relation to phosphorus nutrition. *New Phytologist* 115: 311–317.
- Loveless AR. 1961. A nutritional interpretation of sclerophylly based on differences in the chemical composition of sclerophyllous and mesophytic leaves. *Annals of Botany* 25: 168–184.
- Loveless AR. 1962. Further evidence to support a nutritional interpretation of sclerophylly. *Annals of Botany* 26: 551–561.
- Lurin C, Geelen D, Barbier-Brygoo H, Guern J, Maurel C. 1996. Cloning and functional expression of a plant voltage dependent chloride channel. *Plant Cell* 8: 701–711.
- Lynch J. 1995. Root architecture and plant productivity. *Plant Physiology* 109: 7–13.
- Lynch JP, Brown KM. 2001. Topsoil foraging—an architectural adaptation of plants to low phosphorus. *Plant and Soil* 237: 225–237.
- Ma Z, Bielenberg G, Brown KM, Lynch JP. 2001. Regulation of root hair density by phosphorus availability in *Arabidopsis thaliana*. *Plant, Cell and Environment* 24: 459–467.
- Maathuis FJ, Sanders D. 1999. Plasma membrane transport in context – making sense out of complexity. *Current Opinion in Plant Biology* 2: 236–243.
- Malamy J, Benfey PN. 1997. Organization and cell differentiation in lateral roots of *Arabidopsis thaliana*. *Development* 124: 33–44.
- Marchant A, Bhalerao R, Casimiro I, Eklot J, Casero PJ, Bennett M, Sandberg G. 2002. AUX1 promotes lateral root formation by facilitating indole-3-acetic acid distribution between sink and source tissue in *Arabidopsis* seedling. *Plant Cell* 14: 589–597.
- Marschner H. 1995. *Mineral nutrition of higher plants*, 2nd edn. Boston, MA, USA: Academic Press.
- Marschner H, Romheld V, Cakmak I. 1987. Root-induced changes of nutrient availability in the rhizosphere. *Journal of Plant Nutrition* 10: 1175–1184.
- Marschner H, Römheld V, Horst WJ, Martin P. 1986. Root induced changes in the rhizosphere: importance for mineral nutrition of plants. *Zeitschrift für Pflanzenernährung und Bodenkunde* 149: 441–456.
- Martin AC, del Pozo JC, Iglesias J, Rubio V, Solano R, de la Pena A, Leyva A, Paz-Ares J. 2000. Influence of cytokinins on the expression of phosphate starvation genes in *Arabidopsis*. *Plant Journal* 24: 559–567.
- Masaoka YU, Kojima M, Sugihara S, Yoshihara T, Koshino M, Ichihara A. 1993. Dissolution of ferric phosphate by alfalfa (*Medicago sativa* L.) root exudates. *Plant and Soil* 155/156: 75–78.
- Maser P, Thomine S, Schroeder JI, Ward JM, Hirschi K, Sze H, Talke IN, Amtmann A, Maathuis FJ, Sanders D, Harper JF, Tohieu J, Gribskov M, Persans MW, Salt DE, Kim SA, Guerinst ML. 2001. Phylogenetic relationships within cation transporter families of *Arabidopsis*. *Plant Physiology* 126: 1646–1667.
- Massonneau A, Langlade N, Leon S, Smutny J, Vogt E, Neumann G, Martinoia E. 2001. Metabolic changes associated with cluster root development in white lupin (*Lupinus albus* L.): relationship between organic acid excretion, sucrose metabolism and energy status. *Planta* 213: 534–542.
- Masucci JD, Schiefelbein JW. 1996. Hormones act downstream of TIG and GL2 to promote root hair outgrowth during epidermis development in the *Arabidopsis* root. *Plant Cell* 8: 1505–1517.
- Matar A, Torrent J, Ryan J. 1992. Soil and fertilizer phosphorus and crop responses in the dryland Mediterranean zone. *Advances in Soil Science* 18: 81–146.
- Mathys W. 1977. The role of malate, oxalate, and mustard oil glucosides in the evolution of zinc resistance in herbage plants. *Physiologia Plantarum* 40: 130–136.
- McCulley M. 1995. How do real roots work? *Plant Physiology* 109: 1–6.

- Mertens E. 1991. Pyrophosphate-dependent phosphofructokinase, an anaerobic glycolytic enzyme? *Federation of European Biochemical Societies Letters* 285: 1–5.
- Michael G. 2001. The control of root hair formation: suggested mechanisms. *Journal of Plant Nutrition and Soil Science* 164: 141–145.
- Millar AH, Wiskich JT, Whelan J, Day DA. 1993. Organic acid activation of the alternative oxidase of plant mitochondria. *Federation of European Biochemical Societies Letters* 329: 259–262.
- Miller SS, Liu J, Allan DL, Menzhuber CJ, Fedorova M, Vance CP. 2001. Molecular control of acid phosphatase secretion into the rhizosphere of proteoid roots from phosphorus-stressed white lupin. *Plant Physiology* 127: 594–606.
- Mitsukawa N, Okumura S, Shirano Y, Sato S, Kato T, Harashimi S, Shibata D. 1997. Overexpression of an *Arabidopsis thaliana* high-affinity phosphate transporter gene in tobacco cultured cells enhances cell growth under phosphate-limited conditions. *Proceedings of the National Academy of Sciences, USA* 94: 7098–7102.
- Mok DWS, Mok CM. 2001. Cytokinin metabolism and action. *Annual Review of Plant Physiology and Plant Molecular Biology* 52: 89–118.
- Moraes TF, Plaxton WC. 2000. Purification and characterization of phosphoenolpyruvate carboxylase from *Brassica napus* (rapeseed) suspension cell cultures: implications for phosphoenolpyruvate carboxylase regulation during phosphate starvation, and the integration of glycolysis with nitrogen assimilation. *European Journal of Biochemistry* 267: 4465–4476.
- Morris RO, Bilyeu KD, Laskey JG, Cheikh NN. 1999. Isolation of a gene encoding a glycosylated cytokinin oxidase from maize. *Biochemical and Biophysical Research Communications* 255: 328–333.
- Muchhal US, Pardo JM, Raghothama KG. 1996. Phosphate transporters from the higher plant *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* 93: 10519–10523.
- Muday GK, Haworth P. 1994. Tomato root growth, gravitropism, and lateral development: correlation with auxin transport. *Plant Physiology and Biochemistry* 32: 193–203.
- Murley VR, Theodorou ME, Plaxton WC. 1998. Phosphate starvation-inducible pyrophosphate-dependent phosphofructokinase occurs in plants whose roots do not form symbiotic associations with mycorrhizal fungi. *Physiologia Plantarum* 103: 405–414.
- Nandi SK, Pant RC, Nissen P. 1987. Multiphasic uptake of phosphate by corn roots. *Plant, Cell & Environment* 10: 463–474.
- Narang RA, Bruene A, Altmann T. 2000. Analysis of phosphate acquisition efficiency in different *Arabidopsis* accessions. *Plant Physiology* 124: 1786–1799.
- Neumann G, Martinoia E. 2002. Cluster roots—an underground adaptation for survival in extreme environments. *Trends in Plant Science* 7: 162–167.
- Neumann G, Römheld V. 1999. Root excretion of carboxylic acids and protons in phosphorus-deficient plants. *Plant and Soil* 211: 121–130.
- Neumann G, Massonneau A, Martinoia E, Romheld V. 1999. Physiological adaptations to phosphorus deficiency during proteoid root development in white lupin. *Planta* 208: 373–382.
- Neumann G, Massonneau A, Langlade N, Dinkelaker B, Hengeler C, Romheld V, Martinoia E. 2000. Physiological aspects of cluster root function and development in phosphorus-deficient white lupin (*Lupinus albus* L.). *Annals of Botany* 85: 909–919.
- Ogawa N, Saitoh H, Miura K, Magbanua JP, Bun-ya M, Harashima S, Oshima Y. 1995. Structure and distribution of specific *cis*-elements for transcriptional regulation of PHO84 in *Saccharomyces cerevisiae*. *Molecular and General Genetics* 249: 406–416.
- Oshima Y. 1997. The phosphatase system in *Saccharomyces cerevisiae*. *Genes and Genetic Systems* 72: 323–334.
- Ozawa K, Osaki M, Matsui H, Honma M, Tadano T. 1995. Purification and properties of acid phosphatase secreted from lupin roots under phosphorus deficiency. *Soil Science and Plant Nutrition* 41: 461–469.
- Parker JS, Cavell AC, Dolan L, Roberts K, Grierson CS. 2000. Genetic interactions during root hair morphogenesis in *Arabidopsis*. *Plant Cell* 12: 1961–1974.
- Parsons HL, Yip JY, Vanlerberghe GC. 1999. Increased respiratory restriction during phosphate-limited growth in transgenic tobacco cells lacking alternative oxidase. *Plant Physiology* 121: 1309–1320.
- Pate J, Watt M. 2001. Roots of *Banksia* spp. (Proteaceae) with special reference to functioning of their specialized root clusters. In: Waisel Y, Eshel A, Kafkafi U, eds. *Plant roots: the hidden half*, 3rd edn. New York, NY, USA: Marcel Dekker Inc., 989–1006.
- Pate J, Verboom W, Galloway P. 2001. Co-occurrence of Proteacea, laterite and related oligotrophic soils: coincidental associations or causative inter-relationships? *Australian Journal of Botany* 49: 529–560.
- Peterson RL, Farquhar ML. 1996. Root hairs: specialized tubular cells extending root surfaces. *Botanical Gazette* 62: 1–40.
- Pineros MA, Kochian LV. 2001. A patch-clamp study on the physiology of aluminum toxicity and aluminum tolerance in maize. Identification and characterization of Al(3+)-induced anion channels. *Plant Physiology* 125: 292–305.
- Plaxton WC. 1996. The organization and regulation of plant glycolysis. *Annual Review of Plant Physiology* 47: 185–214.
- Plaxton WC, Carswell MC. 1999. Metabolic aspects of the phosphate starvation response in plants. In: Lerner HR, ed. *Plant responses to environmental stress: from phytohormones to genome reorganization*. New York, NY, USA: Marcel-Dekker, 350–372.
- del Pozo JC, Allona I, Rubio V, Layva A, de la Peña A, Aragoncillo C, Paz-Ares J. 1999. A type 5 acid phosphatase gene from *Arabidopsis thaliana* is induced by phosphate starvation and by some other types of phosphate mobilizing/oxidative stress conditions. *Plant Journal* 19: 579–589.
- Raghothama KG. 1999. Phosphate acquisition. *Annual Review of Plant Physiology and Plant Molecular Biology* 50: 665–693.
- Randall DD, Tolbert NE, Gremel D. 1971. 3-Phosphoglycerate in plants. II. Distribution, physiological considerations, and comparison with phosphoglycolate phosphatase. *Plant Physiology* 48: 480–487.
- Rasmussen AG, Moller IM. 1991. NAD(P)H dehydrogenases on the inner surface of the inner mitochondrial membrane studied using inside-out submitochondrial particles. *Physiologia Plantarum* 83: 357–365.
- Raven JA, Edwards D. 2001. Roots: evolutionary origins and biogeochemical significance. *Journal of Experimental Botany* 52: 381–401.
- Raven JA, Smith FA. 1976. Nitrogen assimilation and transport in vascular land plants in relation to intracellular pH regulation. *New Phytologist* 76: 415–431.
- Reed RC, Brady SR, Muday GK. 1998. Inhibition of auxin movement from the shoot into the root inhibits lateral root development in *Arabidopsis*. *Plant Physiology* 118: 1369–1378.
- Reid JB. 1981. Observations on root hair production by lucerne, maize, and perennial rye grass grown in sandy loam. *Plant and Soil* 62: 319–322.
- Reid RJ, Mimura T, Ohsumi Y, Walker NA, Smith FA. 2000. Phosphate transport in *Chara*: membrane transport via Na/Pi cotransport. *Plant, Cell & Environment* 23: 223–228.
- Richardson AE, Hadobas PA, Hayes JE. 2001. Extracellular secretion of *Aspergillus* phytase from *Arabidopsis* roots enables plants to obtain phosphorus from phytate. *Plant Journal* 25: 1–10.
- Ridge RW. 1995. Recent developments in the cell and molecular biology of root hairs. *Journal of Plant Research* 108: 399–405.
- Rogers E, Guerinot ML. 2002. FRD3, a member of the multidrug and toxin efflux family, controls iron deficiency responses in *Arabidopsis*. *Plant Cell* 14: 1787–1799.
- Rogg LE, Lasswell J, Bartel B. 2001. A gain of function mutation in IAA28 suppresses lateral root development. *Plant Cell* 13: 465–480.
- Romera FJ, Alcantara E. 1994. Iron-deficiency stress response in cucumber (*Cucumis sativus* L.) roots: a possible role for ethylene? *Plant Physiology* 105: 1133–1138.

- Rubio V, Linhares F, Solano R, Martin AC, Iglesias J, Leyva A, Paz-Ares J. 2001. A conserved MYB transcription factor involved in phosphate starvation signaling both in vascular plants and in unicellular algae. *Genes and Development* 15: 2122–2133.
- Ruegger M, Dewey E, Hobbie L, Brown D, Bernasconi P, Turner J, Muday G, Estelle M. 1997. Reduced naphthalamic acid binding in the *tir3* mutant of *Arabidopsis* is associated with a reduction in polar auxin transport and diverse morphological defects. *Plant and Cell* 9: 745–757.
- Runge-Metzger A. 1995. Closing the cycle: obstacles to efficient P management for improved global security. In: Tiessen H, ed. *Phosphorus in the global environment*. Chichester, UK: John Wiley and Sons Ltd, 27–42.
- Russell DW. 1973. *Soil conditions and plant growth*. New York, NY, USA: Longman Group Ltd.
- Ryan PR, Delhaize E, Jones DL. 2001. Function and mechanism of organic anion exudation from plants. *Annual Review of Plant Physiology and Plant Molecular Biology* 52: 527–560.
- Ryan PR, Delhaize E, Randall PJ. 1995a. Characterization of Al-stimulated efflux of malate from the apices of Al-tolerant wheat roots. *Planta* 196: 103–110.
- Ryan PR, Delhaize E, Randall PJ. 1995b. Malate efflux from root apices and tolerance to aluminum are highly correlated in wheat. *Australian Journal of Plant Physiology* 22: 531–536.
- Ryan PR, Skerrett M, Findlay GP, Delhaize E, Tyerman SD. 1997. Aluminum activates an anion channel in the apical cells of wheat roots. *Proceedings of the National Academy of Sciences, USA* 94: 6547–6552.
- Rychter AM, Chauveau M, Bomsel J-L, Lance C. 1992. The effect of phosphate deficiency on mitochondria activity and adenylate levels in bean roots. *Physiologia Plantarum* 84: 80–86.
- Rychter AM, Mikulska M. 1990. The relationship between phosphate status and cyanide-resistant respiration in bean roots. *Physiologia Plantarum* 79: 663–667.
- Sakakibara H, Hayakawa A, Deji A, Gawronski SW, Sugiyama T. 1999. His-Asp phosphotransfer possibly involved in the nitrogen signal transduction mediated by cytokinin in maize: molecular cloning of cDNAs for two-component regulatory factors and demonstration of phosphotransfer activity *in vitro*. *Plant Molecular Biology* 41: 563–573.
- Sakakibara H, Taniguchi M, Sugiyama T. 2000. His-Asp phosphorelay signaling: a communication avenue between plants and their environment. *Plant Molecular Biology* 42: 273–278.
- Sakano K. 2001. Metabolic regulation of pH in plant cells: role of cytoplasmic pH in defense reaction and secondary metabolism. *International Review of Cytology* 206: 1–44.
- Salama AMSE, Wareing PF. 1979. Effect of mineral nutrition on endogenous cytokinins in sunflower (*Helianthus annuus* L.). *Journal of Experimental Botany* 30: 971–981.
- Sanchez PA, Uehara G. 1980. Management consideration for acid soils with high phosphorus fixation capacity. In: Knasawneh FE, Sample EC, Kamrath EJ, eds. *The role of phosphorus in agriculture*. Madison, WI, USA: American Society of Agronomy, 471–514.
- Sanyal SK, DeDatta SK. 1991. Chemistry of phosphorus transformations in soil. *Advances in Soil Science* 16: 1–120.
- Sas L, Rengel Z, Tang C. 2001. Excess cation uptake, and extrusion of protons and organic acid anions by *Lupinus albus* under phosphorus deficiency. *Plant Science* 160: 1191–1198.
- Schachtman DP, Reid RJ, Ayling SM. 1998. Phosphorus uptake by plants: from soil to cell. *Plant Physiology* 116: 447–453.
- Scheres B, Wolkenfelt H. 1998. The *Arabidopsis* root as a model to study plant development. *Plant Physiology and Biochemistry* 36: 21–32.
- Schmidt W. 2001. From faith to fait: ethylene signaling in morphogenic responses to P and Fe deficiency. *Journal of Plant Nutrition and Soil Science* 164: 147–154.
- Schroeder JL. 1995. Anion channels as central mechanisms for signal transduction in guard cells and putative functions in roots for plant–soil interactions. *Plant Molecular Biology* 28: 353–361.
- Shih CY, Kao CH. 1996. Growth inhibition in suspension-cultured rice cells under phosphate deprivation is mediated through putrescine accumulation. *Plant Physiology* 111: 721–724.
- Siedow JN, Umbach AL. 2000. The mitochondrial cyanide-resistant oxidase: structural conservation amid regulatory diversity. *Biochimica et Biophysica Acta* 1459: 432–439.
- Skene KR. 1998. Cluster roots: some ecological considerations. *Journal of Ecology* 86: 1060–1064.
- Skene KR. 2000. Pattern formation in cluster roots: some developmental and evolutionary considerations. *Annals of Botany* 85: 901–908.
- Skene KR, James WM. 2000. A comparison of the effects of auxin on cluster root initiation and development in *Grevillea robusta* Cunn. ex R. Br. (Proteaceae) and the genus *Lupinus* (Leguminosae). *Plant and Soil* 219: 221–229.
- Skene KR, Kierans M, Sprent J, Raven JA. 1996. Structural aspects of cluster root development and their possible significance for nutrient acquisition in *Grevillea robusta* (Proteaceae). *Annals of Botany* 77: 443–451.
- Skoog F, Miller CO. 1957. Chemical regulation of growth and organ formation in plant tissue culture *in vitro*. *Symposium of the Society of Experimental Biology* 11: 118–131.
- Smith FA, Jakobsen I, Smith SE. 2000. Spatial differences in acquisition of soil phosphate between two arbuscular mycorrhizal fungi in symbiosis with *Medicago truncatula*. *New Phytologist* 147: 357–366.
- Smith FW. 2001. Sulphur and phosphorus transport systems in plants. *Plant Soil* 232: 109–118.
- Smith FW, Ealing PM, Dong B, Delhaize E. 1997. The cloning of two *Arabidopsis* genes belonging to a phosphate transporter family. *Plant Journal* 11: 83–92.
- Smith FW, Rae AL, Hawkesford MJ. 2000. Molecular mechanisms of phosphate and sulfate transport in plants. *Biochimica et Biophysica Acta* 1465: 236–245.
- Smith SE, Dickson S, Smith FA. 2001. Nutrient transfer in arbuscular mycorrhizas: how are fungal and plant processes integrated? *Australian Journal of Plant Physiology* 28: 683–694.
- Tadano T, Sakai H. 1991. Secretion of acid phosphatase by roots of several crop species under phosphorus-deficient conditions. *Soil Science and Plant Nutrition* 37: 129–140.
- Takahashi A, Takeda K, Ohnishi T. 1991. Light-induced anthocyanin reduces the extent of damage to DNA in UV-irradiated *Centaurea cyanus* cells in culture. *Plant and Cell Physiology* 32: 541–547.
- Tang Z, Sadka A, Morishige DT, Mullet JE. 2001. Homeodomain leucine zipper proteins bind to the phosphate response domain of the soybean *VspB* tripartite promoter. *Plant Physiology* 125: 797–809.
- Tanimoto M, Roberts K, Dolan L. 1995. Ethylene is a positive regulator of root hair development in *Arabidopsis thaliana*. *Plant Journal* 8: 943–948.
- Tarafdar JC, Claassen N. 1988. Organic phosphorus compounds as a phosphorus source for higher plants through the activity of phosphatases produced by plant roots and microorganisms. *Biology and Fertility of Soils* 5: 3308–3312.
- Tesfaye M, Temple SJ, Allan DL, Vance CP, Samac DA. 2001. Overexpression of malate dehydrogenase in transgenic alfalfa enhances organic acid synthesis and confers tolerance to aluminum. *Plant Physiology* 127: 1836–1844.
- Theodorou ME, Cornel FA, Duff SM, Plaxton WC. 1992. Phosphate starvation-inducible synthesis of the alpha-subunit of the pyrophosphate-dependent phosphofructokinase in black mustard suspension cells. *Journal of Biological Chemistry* 267: 21901–21905.
- Theodorou ME, Plaxton WC. 1993. Metabolic adaptations of plant respiration to nutritional phosphate deprivation. *Plant Physiology* 101: 339–344.
- Theodorou ME, Plaxton WC. 1996. Purification and characterization of pyrophosphate-dependent phosphofructokinase from phosphate-starved *Brassica nigra* suspension cells. *Plant Physiology* 112: 343–351.
- Theodoulou FL. 2000. Plant ABC transporters. *Biochimica et Biophysica Acta* 1465: 79–103.

- Tibbett M, Sanders FE. 2002. Ectomycorrhizal symbiosis can enhance plant nutrition through improved access to discrete organic nutrient patches of high resource quality. *Annals of Botany* 89: 783–789.
- Tillberg J-E, Rowley JR. 1989. Physiological and structural effects of phosphorus starvation on the unicellular green alga *Scenedesmus*. *Physiologia Plantarum* 75: 315–324.
- Torrey JG. 1986. Endogenous and exogenous influences in the regulation of lateral root formation. In: Jackson AB, ed. *New root formation in plants and cuttings*. Dordrecht, The Netherlands: Martinus-Nijhoff.
- Turner IM. 1994. Sclerophylly: primarily protective? *Functional Ecology* 8: 669–675.
- von Uexküll HR, Mutert E. 1995. Global extent, development and economic impact of acid soils. *Plant and Soil* 171: 1–15.
- Uhde-Stone C, Gilbert G, Johnson JMF, Litjens R, Zinn KE, Temple SJ, Vance CP, Allan DL. 2003a. Adaptation of white lupin to phosphorus deficiency involves enhanced expression of genes related to organic acid metabolism. *Plant and Soil* (In press.)
- Uhde-Stone C, Zinn KE, Ramirez-Yáñez M, Li A, Vance CP, Allan DL. 2003b. Nylon filter arrays reveal differential gene expression in proteoid roots of white lupin in response to P deficiency. *Plant Physiology* 131 (In press.)
- Ullrich C, Nováček A. 1990. Extra- and intracellular pH and membrane potential changes induced by K^+ , Cl^- , $H_2PO_4^-$ and NO_3^- uptake and fusicoccin in root hairs of *Limnobium stoloniferum*. *Plant Physiology* 94: 1561–1567.
- Urao T, Yakubov B, Satoh R, Yamaguchi-Shinozaki K, Seki M, Hirayama T, Shinozaki K. 1999. A transmembrane hybrid-type histidine kinase in *Arabidopsis* functions as an osmosensor. *Plant Cell* 11: 1743–1754.
- Urao T, Yamaguchi-Shinozaki K, Shinozaki K. 2000. Two-component system in plant signal transduction. *Trends in Plant Science* 5: 67–74.
- Vance CP. 2001. Symbiotic nitrogen fixation and phosphorus acquisition: plant nutrition in a world of declining renewable resources. *Plant Physiology* 127: 390–397.
- Vance CP, Lamb JFS. 2001. Application of biochemical studies to improving nitrogen fixation. *Australian Journal of Experimental Agriculture* 41: 403–416.
- Vogel K, Hinnen A. 1990. The yeast phosphatase system. *Molecular Microbiology* 4: 2013–2017.
- de Vos CR, Lubberding HJ, Biefait HF. 1986. Rhizosphere acidification as a response to iron deficiency in bean plants. *Plant Physiology* 81: 842–846.
- Wagner B, Beck F. 1993. Cytokinins in the perennial herb *Urtica dioica* L. as influenced by nitrogen status. *Planta* 190: 511–518.
- Wanner BL. 1993. Gene regulation by phosphate in enteric bacteria. *Journal of Cell Biochemistry* 51: 47–54.
- Wasaki J, Ando M, Ozawa K, Omura M, Osaki M, Ito H, Matsui H, Tadano T. 1997. Properties of secretory acid phosphatase from lupin roots under phosphorus-deficient conditions. *Soil Science and Plant Nutrition* 43: 981–986.
- Wasaki J, Omura M, Osaki M, Ito H, Matsui H, Shinano T, Tadano T. 1999. Structure of a cDNA for an acid phosphatase from phosphate-deficient lupin (*Lupinus albus* L.) roots. *Soil Science and Plant Nutrition* 45: 439–449.
- Wasaki J, Yamamura T, Shinano T, Osaki M. 2003. Secreted acid phosphatase is expressed in cluster roots of lupin in response to phosphorus deficiency. *Plant and Soil* (In press.)
- Waters BM, Blevins DG. 2000. Ethylene production, cluster root formation, and localization of iron (III) reducing capacity in Fe deficient squash roots. *Plant and Soil* 225: 21–31.
- Watt M, Evans JR. 1999. Proteoid roots. Physiology and development. *Plant Physiology* 121: 317–324.
- Weger HG, Dasgupta R. 1993. Regulation of alternative pathway respiration in *Chlamydomonas reinhardtii* (Chlorophyceae). *Journal of Phycology* 29: 300–308.
- Williamson LC, Ribrioux SP, Fitter AH, Leyser HM. 2001. Phosphate availability regulates root system architecture in *Arabidopsis*. *Plant Physiology* 126: 875–882.
- Wissuwa M, Ae N. 2001. Further characterization of two QTLs that increase phosphorus uptake of rice (*Oryza sativa* L.) under phosphorus deficiency. *Plant and Soil* 237: 275–286.
- Wykoff DD, Grossman AR, Weeks DP, Usuda H, Shimogawara K. 1999. Psr1, a nuclear localized protein that regulates phosphorus metabolism in *Chlamydomonas*. *Proceedings of the National Academy of Sciences, USA* 96: 15336–15341.
- Yu B, Xu C, Benning C. 2002. *Arabidopsis* disrupted in *SQD2* encoding sulfolipid synthase is impaired in phosphate-limited growth. *National Academy of Sciences, USA* 99: 5732–5737.
- Yamakawa T, Kato S, Ishida K, Kodama T, Minoda Y. 1983. Production of anthocyanin by *Vitis* cell in suspension culture. *Agricultural and Biological Chemistry* 47: 2185–2191.
- Yan X. 1999. Phosphorus efficiency of cultivated legumes in agroecosystems: the case of South China. In: Lynch JP, Deikman J, eds. *Phosphorus in plant biology: regulatory roles in molecular, cellular, organismic, and ecosystem processes*. Rockville, MD, USA: American Society of Plant Physiology, 85–93.
- Yan F, Zhu Y, Mueller C, Schubert S. 2002. Adaptation of H^+ pumping and plasma membrane H^+ ATPase activity in proteoid roots of white lupin under phosphate deficiency. *Plant Physiology* 129: 50–63.
- Yang Y-Y, Jung J-Y, Song W-Y, Suh H-S, Lee Y. 2000. Identification of rice varieties with high tolerance or sensitivity to lead and characterization of the mechanism of tolerance. *Plant Physiology* 124: 1019–1026.
- Zhang H, Forde BG. 1998. An *Arabidopsis* MADS box gene that controls nutrient-induced changes in root architecture. *Science* 279: 407–409.
- Zhang H, Forde BG. 2000. Regulation of *Arabidopsis* root development by nitrate availability. *Journal of Experimental Botany* 342: 51–59.
- Zhang C, McManus M. 2000. Identification and characterization of two distinct acid phosphatases in cell walls of white clover. *Plant Physiology and Biochemistry* 38: 259–270.
- Zhang WH, Ryan PR, Tyerman SD. 2001. Malate-permeable channels and cation channels activated by aluminum in the apical cells of wheat roots. *Plant Physiology* 125: 1459–1472.